

Non-Equilibrium Dynamics of Dyson's Model with an Infinite Number of Particles

Makoto Katori¹, Hideki Tanemura²

- 1 Department of Physics, Faculty of Science and Engineering, Chuo University, Kasuga, Bunkyo-ku, Tokyo 112-8551, Japan. E-mail: katori@phys.chuo-u.ac.jp
- 2 Department of Mathematics and Informatics, Faculty of Science, Chiba University, 1-33 Yayoi-cho, Inage-ku, Chiba 263-8522, Japan. E-mail: tanemura@math.s.chiba-u.ac.jp

(19 June 2009)

Abstract: Dyson's model is a one-dimensional system of Brownian motions with long-range repulsive forces acting between any pair of particles with strength proportional to the inverse of distances with proportionality constant $\beta/2$. We give sufficient conditions for initial configurations so that Dyson's model with $\beta = 2$ and an infinite number of particles is well defined in the sense that any multitime correlation function is given by a determinant with a continuous kernel. The class of infinite-dimensional configurations satisfying our conditions is large enough to study non-equilibrium dynamics. For example, we obtain the relaxation process starting from a configuration, in which every point of $\mathbb{Z}$ is occupied by one particle, to the stationary state, which is the determinantal point process with the sine kernel.

1 Introduction

In order to understand the statistics of eigenvalues of random matrix ensembles as equilibrium distributions of particle positions in the one-dimensional Coulomb gas systems with log-potentials, Dyson introduced stochastic models of particles in $\mathbb{R}$, which obey the stochastic differential equations (SDEs),

$$dX_j(t) = dB_j(t) + \frac{\beta}{2} \sum_{1 \leq k \leq N, k \neq j} \frac{dt}{X_j(t) - X_k(t)}, \quad 1 \leq j \leq N, \quad t \in [0, \infty), \quad (1.1)$$

where $B_j(t)$'s are independent one-dimensional standard Brownian motions [3]. The Gaussian orthogonal ensemble (GOE), the Gaussian unitary ensemble (GUE), and the Gaussian symplectic ensemble (GSE) of random matrices correspond to the SDEs (1.1) with $\beta = 1, 2$ and 4, respectively [14]. Spohn [20] has considered an infinite particle system obtained by taking the $N \rightarrow \infty$ limit of (1.1) with $\beta = 2$ and called the system *Dyson's model*. He studied the equilibrium dynamics with respect to the determinantal (Fermion) point process $\mu_{\sin}$, in which any spatial correlation function ρ_m is given by a determinant with the *sine kernel* [19, 18]

$$K_{\sin}(y - x) = \frac{1}{2\pi} \int_{|k| \leq \pi} dk e^{ik(y-x)} = \frac{\sin\{\pi(y-x)\}}{\pi(y-x)}, \quad x, y \in \mathbb{R}, \quad (1.2)$$

where $i = \sqrt{-1}$. By the Dirichlet form approach Osada [16] constructed the infinite particle system represented by a diffusion process, which has μ_{sin} as a reversible measure. Recently he proved that this system satisfies the SDEs (1.1) with $N = \infty$ [17]. On the other hand, it was shown by Eynard and Mehta [4] that multitime correlation functions for the process (1.1) are generally given by determinants, if the process starts from $\mu_{N,\sigma^2}^{\text{GUE}}$, the eigenvalue distribution of GUE with variance σ^2 . Nagao and Forrester [15] evaluated the bulk scaling limit $\sigma^2 = 2N/\pi^2 \rightarrow \infty$ and derived the so-called *extended sine kernel* with density 1,

$$\begin{aligned} \mathbf{K}_{\text{sin}}(t-s, y-x) &= \frac{1}{2\pi} \int_{|k| \leq \pi} dk e^{k^2(t-s)/2 + ik(y-x)} - \mathbf{1}(s > t)p(s-t, x|y) \\ &= \begin{cases} \int_0^1 du e^{\pi^2 u^2(t-s)/2} \cos\{\pi u(y-x)\} & \text{if } t > s \\ K_{\text{sin}}(y-x) & \text{if } t = s \\ - \int_1^\infty du e^{\pi^2 u^2(t-s)/2} \cos\{\pi u(y-x)\} & \text{if } t < s, \end{cases} \quad (1.3) \end{aligned}$$

$s, t \geq 0, x, y \in \mathbb{R}$, where $\mathbf{1}(\omega)$ is the indicator function of condition ω , and $p(t, y|x)$ is the *heat kernel*

$$p(t, y|x) = \frac{e^{-(y-x)^2/2t}}{\sqrt{2\pi t}} = \frac{1}{2\pi} \int_{\mathbb{R}} dk e^{-k^2 t/2 + ik(y-x)}, \quad t > 0. \quad (1.4)$$

Since $\lim_{N \rightarrow \infty} \mu_{N, 2N/\pi^2}^{\text{GUE}} = \mu_{\text{sin}}$, the process, whose multitime correlation functions are given by determinants with the extended sine kernel (1.3), is expected to be identified with the infinite-dimensional equilibrium dynamics of Spohn and Osada. This equivalence is, however, not yet proved.

Fritz [5] established the theory of non-equilibrium dynamics of infinite particle systems with a finite-range smooth potential. Here we study the non-equilibrium dynamics of infinite-particle Dyson's model with a long-range log-potential, in which the force acting each particle is singular both for short and long distances (see (1.1)).

We denote by $\mathfrak{M}$ the space of nonnegative integer-valued Radon measures on $\mathbb{R}$, which is a Polish space with the vague topology: we say $\xi_n, n \in \mathbb{N} \equiv \{1, 2, \dots\}$ converges to ξ vaguely, if $\lim_{n \rightarrow \infty} \int_{\mathbb{R}} \varphi(x) \xi_n(dx) = \int_{\mathbb{R}} \varphi(x) \xi(dx)$ for any $\varphi \in C_0(\mathbb{R})$, where $C_0(\mathbb{R})$ is the set of all continuous real-valued functions with compact supports. Any element ξ of $\mathfrak{M}$ can be represented as $\xi(\cdot) = \sum_{j \in \Lambda} \delta_{x_j}(\cdot)$ with a sequence of points in $\mathbb{R}$, $\mathbf{x} = (x_j)_{j \in \Lambda}$ satisfying $\xi(K) = \#\{j \in \Lambda : x_j \in K\} < \infty$ for any compact subset $K \subset \mathbb{R}$. The index set Λ is $\mathbb{N}$ or a finite set. We call an element ξ of $\mathfrak{M}$ an unlabeled configuration, and a sequence $\mathbf{x}$ a labeled configuration. For $A \subset \mathbb{R}$, we write the restriction of ξ on A as $(\xi \cap A)(\cdot) = \sum_{j \in \Lambda: x_j \in A} \delta_{x_j}(\cdot)$.

As an $\mathfrak{M}$ -valued process $(\mathbb{P}, \Xi(t), t \in [0, \infty))$, we consider the system such that, for any integer $M \geq 1$, $f_m \in C_0(\mathbb{R}), \theta_m \in \mathbb{R}, 1 \leq m \leq M, 0 < t_1 < \dots <$

$t_M < \infty$, the expectation of $\exp \left\{ \sum_{m=1}^M \theta_m \int_{\mathbb{R}} f_m(x) \Xi(t_m, dx) \right\}$ can be expanded with $\chi_m(x) = e^{\theta_m f_m(x)} - 1, 1 \leq m \leq M$ as

$$\begin{aligned} \mathcal{G}^\xi[\chi] \equiv & \sum_{N_1 \geq 0} \cdots \sum_{N_M \geq 0} \prod_{m=1}^M \frac{1}{N_m!} \int_{\mathbb{R}^{N_1}} \prod_{j=1}^{N_1} dx_j^{(1)} \cdots \int_{\mathbb{R}^{N_M}} \prod_{j=1}^{N_M} dx_j^{(M)} \\ & \times \prod_{m=1}^M \prod_{j=1}^{N_m} \chi_m(x_j^{(m)}) \rho(t_1, \mathbf{x}_{N_1}^{(1)}; \dots; t_M, \mathbf{x}_{N_M}^{(M)}), \end{aligned}$$

where $\mathbf{x}_{N_m}^{(m)}$ denotes $(x_1^{(m)}, \dots, x_{N_m}^{(m)})$, $1 \leq m \leq M$. Here ρ 's are locally integrable functions, which are symmetric in the sense that

$$\rho(\dots; t_m, \sigma(\mathbf{x}_{N_m}^{(m)}); \dots) = \rho(\dots; t_m, \mathbf{x}_{N_m}^{(m)}; \dots) \quad \text{with} \quad \sigma(\mathbf{x}_{N_m}^{(m)}) \equiv (x_{\sigma(1)}^{(m)}, \dots, x_{\sigma(N_m)}^{(m)})$$

for any permutation $\sigma \in \mathcal{S}_{N_m}, 1 \leq \forall m \leq M$. In such a system $\rho(t_1, \mathbf{x}_{N_1}^{(1)}; \dots; t_M, \mathbf{x}_{N_M}^{(M)})$ is called the $(N_1, \dots, N_M)$ -multitime correlation function and $\mathcal{G}^\xi[\chi]$ the generating function of multitime correlation functions. There are no multiple points with probability one for $t > 0$. Then we assume that there is a function $\mathbb{K}(s, x; t, y)$, which is continuous with respect to $(x, y) \in \mathbb{R}^2$ for any fixed $(s, t) \in [0, \infty)^2$, such that

$$\rho(t_1, \mathbf{x}_{N_1}^{(1)}; \dots; t_M, \mathbf{x}_{N_M}^{(M)}) = \det_{\substack{1 \leq j \leq N_m, 1 \leq k \leq N_n \\ 1 \leq m, n \leq M}} \left[\mathbb{K}(t_m, x_j^{(m)}; t_n, x_k^{(n)}) \right]$$

for any integer $M \geq 1$, any sequence $(N_m)_{m=1}^M$ of positive integers, and any time sequence $0 < t_1 < \dots < t_M < \infty$. That is, the finite dimensional distributions of the process are determined by the function $\mathbb{K}$. Let $\mathbf{T} = \{t_1, \dots, t_M\}$. We note that $\Xi^{\mathbf{T}} = \sum_{t \in \mathbf{T}} \delta_t \otimes \Xi(t)$ is a determinantal (Fermion) point process on $\mathbf{T} \times \mathbb{R}$ with an operator $\mathcal{K}$ given by $\mathcal{K}f(s, x) = \sum_{t \in \mathbf{T}} \int_{\mathbb{R}} dy \mathbb{K}(s, x; t, y) f(t, y)$ for $f(t, \cdot) \in C_0(\mathbb{R}), t \in \mathbf{T}$. The process is then said to be *determinantal with the correlation kernel* $\mathbb{K}$. When $\mathcal{K}$ is symmetric, Soshnikov [19] and Shirai and Takahashi [18] gave sufficient conditions for $\mathbb{K}$ to be a correlation kernel of a determinantal point process. Though such conditions are not known for asymmetric cases, a variety of processes, which are determinantal with asymmetric correlation kernels, have been studied. As mentioned above the process $\Xi(t) = \sum_{j=1}^N \delta_{X_j(t)}$ with the SDEs (1.1) with $\beta = 2$ starting from its equilibrium measure $\mu_{N, \sigma^2}^{\text{GUE}}$ is an example [4]. The infinite particle system of Nagao and Forrester [15] is also determinantal with the extended sine kernel, which is asymmetric as shown by (1.3). (For other examples, see, for instance, [21, 10].)

In the present paper we first show that, for any fixed configuration $\xi^N \in \mathfrak{M}$ with $\xi^N(\mathbb{R}) = N$, Dyson's model starting from ξ^N is determinantal and its correlation kernel $\mathbb{K}^{\xi^N}$ is given by using the *multiple Hermite polynomials* [8, 2, 7] (Proposition 2.1).

For $\xi \in \mathfrak{M}$, when $\mathbb{K}^{\xi \cap [-L, L]}$ converges to a continuous function as $L \rightarrow \infty$, the limit is written as $\mathbb{K}^\xi$. If $\mathbb{P}_{\xi \cap [-L, L]}$ converges to a probability measure $\mathbb{P}_\xi$ on $\mathfrak{M}^{[0, \infty)}$, which is determinantal with the correlation kernel $\mathbb{K}^\xi$, weakly in the sense of finite dimensional distributions as $L \rightarrow \infty$ in the vague topology, we say that the process $(\mathbb{P}_\xi, \Xi(t), t \in [0, \infty))$ is *well defined with the correlation kernel* $\mathbb{K}^\xi$. (The regularity of the sample paths of $\Xi(t)$ will be discussed elsewhere [11].) In the case $\xi(\mathbb{R}) = \infty$, the process $(\mathbb{P}_\xi, \Xi(t), t \in [0, \infty))$ is Dyson's model with an infinite number of particles.

For $\xi \in \mathfrak{M}$ with $\xi(\{x\}) \leq 1, \forall x \in \mathbb{R}$, we give sufficient conditions so that the process $(\mathbb{P}_\xi, \Xi(t), t \in [0, \infty))$ is well defined, in which the correlation kernel is generally expressed using a double integral with the heat kernels of an *entire function* represented by an infinite product (Theorem 2.2). The configuration in which every point of $\mathbb{Z}$ is occupied by one particle, $\xi^\mathbb{Z}(\cdot) \equiv \sum_{\ell \in \mathbb{Z}} \delta_\ell(\cdot)$, satisfies the conditions and we will show that Dyson's model starting from $\xi^\mathbb{Z}$ is determinantal with the kernel

$$\begin{aligned} \mathbb{K}^{\xi^\mathbb{Z}}(s, x; t, y) &= \mathbf{K}_{\sin}(t - s, y - x) \\ &\quad + \frac{1}{2\pi} \int_{|k| \leq \pi} dk e^{k^2(t-s)/2 + ik(y-x)} \left\{ \vartheta_3(x - iks, 2\pi is) - 1 \right\} \\ &= \mathbf{K}_{\sin}(t - s, y - x) \\ &\quad + \sum_{\ell \in \mathbb{Z} \setminus \{0\}} e^{2\pi i x \ell - 2\pi^2 s \ell^2} \int_0^1 du e^{\pi^2 u^2(t-s)/2} \cos \left[\pi u \{ (y - x) - 2\pi i s \ell \} \right], \end{aligned} \quad (1.5)$$

$s, t \geq 0, x, y \in \mathbb{R}$, where ϑ_3 is a version of the Jacobi theta function defined by

$$\vartheta_3(v, \tau) = \sum_{\ell \in \mathbb{Z}} e^{2\pi i v \ell + \pi i \tau \ell^2}, \quad \Im \tau > 0. \quad (1.6)$$

The lattice structure $\mathbb{K}^{\xi^\mathbb{Z}}(s, x + n; t, y + n) = \mathbb{K}^{\xi^\mathbb{Z}}(s, x; t, y), \forall n \in \mathbb{Z}, s, t \geq 0$ is clear in (1.5) by the periodicity of ϑ_3 , $\vartheta_3(v + n, \tau) = \vartheta_3(v, \tau), \forall n \in \mathbb{Z}$. We can prove

$$\lim_{u \rightarrow \infty} \mathbb{K}^{\xi^\mathbb{Z}}(u + s, x; u + t, y) = \mathbf{K}_{\sin}(t - s, y - x), \quad (1.7)$$

which implies that $\mu_{\sin}$ is an attractor of Dyson's model and $\xi^\mathbb{Z}$ is in its basin.

We are interested in the continuity of the process with respect to initial configuration. For Dyson's model with finite particles, the weak convergence of the processes $(\mathbb{P}_{\xi_n^N}, \Xi(t), t \in [0, \infty)) \rightarrow (\mathbb{P}_{\xi^N}, \Xi(t), t \in [0, \infty))$ as $n \rightarrow \infty$ is guaranteed by the vague convergence of the initial configurations $\xi_n^N \rightarrow \xi^N$ as $n \rightarrow \infty$, where $\xi_n^N, \xi^N \in \mathfrak{M}$ with $\xi_n^N(\mathbb{R}) = \xi^N(\mathbb{R}) = N < \infty, n \in \mathbb{N}$. Based on this continuity, Dyson's model can be defined for any initial configurations with finite particles, which can have multiple points (see Proposition 2.1). On the other hand, we have found that, if $\xi(\mathbb{R}) = \infty$, the weak convergence of processes in the sense of finite dimensional distributions cannot be concluded from the convergence of initial configurations in the vague topology. In the present paper we consider a stronger topology for infinite-particle

configurations (Definition 2.3). We introduce the spaces $\mathfrak{Y}_m^\kappa, \kappa \in (1/2, 1), m \in \mathbb{N}$ of initial configurations such that the convergence of processes is guaranteed by that of the initial configurations in this new topology (Theorem 2.4).

Note that the union of the spaces $\mathfrak{Y} = \bigcup_{\kappa \in (1/2, 1)} \bigcup_{m \in \mathbb{N}} \mathfrak{Y}_m^\kappa$ is large enough to carry the Poisson point processes, Gibbs states with regular conditions, μ_{sin} , as well as infinite-particle configurations with multiple points. In particular, using the fact $\mu_{\text{sin}}(\mathfrak{Y}) = 1$ and the continuity with respect to the initial configurations, we can prove that the process $(\mathbf{P}_{\text{sin}}, \Xi(t), t \in [0, \infty))$ of Nagao and Forrester, which is determinantal with the extended sine kernel (1.3), is Markovian [11].

The paper is organized as follows. In Section 2 preliminaries and main results are given. In Section 3 the definitions of some special functions used in the present paper are given and their basic properties are summarized. Section 4 is devoted to proofs of results.

2 Preliminaries and Main Results

For $\xi(\cdot) = \sum_{j \in \Lambda} \delta_{x_j}(\cdot) \in \mathfrak{M}$, we introduce the following operations;

(shift) for $u \in \mathbb{R}$, $\tau_u \xi(\cdot) = \sum_{j \in \Lambda} \delta_{x_j + u}(\cdot)$,

(dilatation) for $c > 0$, $c \circ \xi(\cdot) = \sum_{j \in \Lambda} \delta_{cx_j}(\cdot)$,

(square) $\xi^{(2)}(\cdot) = \sum_{j \in \Lambda} \delta_{x_j^2}(\cdot)$.

We use the convention such that

$$\prod_{x \in \xi} f(x) = \exp \left\{ \int_{\mathbb{R}} \xi(dx) \log f(x) \right\} = \prod_{x \in \text{supp } \xi} f(x)^{\xi(\{x\})}$$

for $\xi \in \mathfrak{M}$ and a function f on $\mathbb{R}$, where $\text{supp } \xi = \{x \in \mathbb{R} : \xi(\{x\}) > 0\}$. For a multivariate symmetric function g we write $g((x)_{x \in \xi})$ for $g((x_j)_{j \in \Lambda})$.

For $s, t \in [0, \infty)$, $x, y \in \mathbb{R}$ and $\xi^N \in \mathfrak{M}$ with $\xi^N(\mathbb{R}) = N \in \mathbb{N}$, we set

$$\begin{aligned} \mathbb{K}^{\xi^N}(s, x; t, y) &= \frac{1}{2\pi i} \oint_{\Gamma(\xi^N)} dz p(s, x|z) \int_{\mathbb{R}} dy' p(t, -iy|y') \\ &\quad \times \frac{1}{iy' - z} \prod_{x' \in \xi^N} \left(1 - \frac{iy' - z}{x' - z} \right) \\ &\quad - \mathbf{1}(s > t) p(s - t, x|y), \end{aligned} \tag{2.1}$$

where $\Gamma(\xi^N)$ is a closed contour on the complex plane $\mathbb{C}$ encircling the points in $\text{supp } \xi^N$ on the real line $\mathbb{R}$ once in the positive direction.

Proposition 2.1 *Dyson's model $(\mathbb{P}_{\xi^N}, \Xi(t), t \in [0, \infty))$, starting from any fixed configuration $\xi^N \in \mathfrak{M}$ with $\xi^N(\mathbb{R}) = N < \infty$, is determinantal with the correlation kernel $\mathbb{K}^{\xi^N}$ given by (2.1).*

We put

$$\mathfrak{M}_0 = \left\{ \xi \in \mathfrak{M} : \xi(\{x\}) \leq 1 \text{ for any } x \in \mathbb{R} \right\}.$$

Since any element ξ of $\mathfrak{M}_0$ is determined uniquely by its support, it is identified with a countable subset $\{x_j\}_{j \in \Lambda}$ of $\mathbb{R}$. For $\xi^N \in \mathfrak{M}_0$, $a \in \mathbb{C}$, we introduce an entire function of $z \in \mathbb{C}$

$$\Phi(\xi^N, a, z) = \prod_{x \in \xi^N \cap \{a\}^c} \left(1 - \frac{z - a}{x - a} \right),$$

whose zero set is $\text{supp } (\xi^N \cap \{a\}^c)$ (see, for instance, [12]). Then, if $\xi^N \in \mathfrak{M}_0$, (2.1) is written as

$$\begin{aligned} \mathbb{K}^{\xi^N}(s, x; t, y) &= \int_{\mathbb{R}} \xi^N(dx') p(s, x|x') \int_{\mathbb{R}} dy' p(t, -iy|y') \Phi(\xi^N, x', iy') \\ &\quad - \mathbf{1}(s > t) p(s - t, x|y). \end{aligned} \quad (2.2)$$

For $L > 0$, $\alpha > 0$ and $\xi \in \mathfrak{M}$ we put

$$M(\xi, L) = \int_{[-L, L] \setminus \{0\}} \frac{\xi(dx)}{x}, \quad M_\alpha(\xi, L) = \left(\int_{[-L, L] \setminus \{0\}} \frac{\xi(dx)}{|x|^\alpha} \right)^{1/\alpha},$$

and

$$M(\xi) = \lim_{L \rightarrow \infty} M(\xi, L), \quad M_\alpha(\xi) = \lim_{L \rightarrow \infty} M_\alpha(\xi, L),$$

if the limits finitely exist. We introduce the following conditions:

(C.1) there exists $C_0 > 0$ such that $|M(\xi)| < C_0$,

(C.2) (i) there exist $\alpha \in (1, 2)$ and $C_1 > 0$ such that $M_\alpha(\xi) \leq C_1$,

(ii) there exist $\beta > 0$ and $C_2 > 0$ such that

$$M_1(\tau_{-a^2} \xi^{(2)}) \leq C_2(|a| \vee 1)^{-\beta} \quad \forall a \in \text{supp } \xi.$$

We denote by $\mathfrak{X}$ the set of configurations ξ satisfying the conditions (C.1) and (C.2), and put $\mathfrak{X}_0 = \mathfrak{X} \cap \mathfrak{M}_0$. For $\xi \in \mathfrak{X}_0$, $a \in \mathbb{R}$ and $z \in \mathbb{C}$ we define

$$\Phi(\xi, a, z) = \lim_{L \rightarrow \infty} \Phi(\xi \cap [a - L, a + L], a, z).$$

We note that $|\Phi(\xi, a, z)| < \infty$ and $\Phi(\xi, a, \cdot) \not\equiv 0$, if $|M(\tau_{-a} \xi)| < \infty$ and $M_2(\tau_{-a} \xi) < \infty$.

Theorem 2.2 *If $\xi \in \mathfrak{X}_0$, the process $(\mathbb{P}_\xi, \Xi(t), t \in [0, \infty))$ is well defined with the correlation kernel given by*

$$\begin{aligned} \mathbb{K}^\xi(s, x; t, y) &= \int_{\mathbb{R}} \xi(dx') p(s, x|x') \int_{\mathbb{R}} dy' p(t, -iy|y') \Phi(\xi, x', iy') \\ &\quad - \mathbf{1}(s > t) p(s - t, x|y). \end{aligned} \quad (2.3)$$

In case $\xi(\mathbb{R}) = \infty$, Theorem 2.2 gives Dyson's model with an infinite number of particles starting from the configuration $\xi \in \mathfrak{X}_0$. From (2.3) it is easy to check that

$$\mathbb{K}^\xi(t, x; t, y) \mathbb{K}^\xi(t, y; t, x) dx dy \rightarrow \xi(dx) \mathbf{1}(x = y), \quad t \rightarrow 0 \quad \text{in the vague topology.}$$

An interesting and important example is obtained for the initial configuration, in which every point in $\mathbb{Z}$ is occupied by one particle, $\xi^\mathbb{Z}(\cdot) \equiv \sum_{\ell \in \mathbb{Z}} \delta_\ell(\cdot)$. In this case $\xi^\mathbb{Z}(\cdot) \in \mathfrak{X}_0$ and we can show that the correlation kernel $\mathbb{K}^{\xi^\mathbb{Z}}$ is given by (1.5). The process $(\mathbf{P}_{\text{sin}}, \Xi(t), t \in [0, \infty))$ is reversible with respect to μ_{sin} . The result (1.7) implies that the process $(\mathbb{P}_{\xi^\mathbb{Z}}, \Xi(u+t), t \in [0, \infty))$ converges to $(\mathbf{P}_{\text{sin}}, \Xi(t), t \in [0, \infty))$, as $u \rightarrow \infty$, weakly in the sense of finite dimensional distributions. In other words, $(\mathbb{P}_{\xi^\mathbb{Z}}, \Xi(t), t \in [0, \infty))$ is the *relaxation process* from an initial configuration $\xi^\mathbb{Z}$ to the invariant measure μ_{sin} , which is determinantal, and this non-equilibrium dynamics is completely determined via the temporally inhomogeneous correlation kernel (1.5). (See Remark in Section 4.3.)

For $\kappa > 0$, we put

$$g^\kappa(x) = \text{sgn}(x)|x|^\kappa, \quad x \in \mathbb{R}, \quad \text{and} \quad \eta^\kappa(\cdot) = \sum_{\ell \in \mathbb{Z}} \delta_{g^\kappa(\ell)}(\cdot).$$

Since g^κ is an odd function, η^κ satisfies (C.1) for any $\kappa > 0$. For any $\kappa > 1/2$ we can show by simple calculation that η^κ satisfies (C.2)(i) with any $\alpha \in (1/\kappa, 2)$ and some $C_1 = C_1(\alpha) > 0$ depending on α , and does (C.2)(ii) with any $\beta \in (0, 2\kappa - 1)$ and some $C_2 = C_2(\beta) > 0$ depending on β . This implies that η^κ is an element of $\mathfrak{X}_0$ in any case $\kappa > 1/2$. Note that $\eta^1 = \xi^\mathbb{Z}$.

If there exists $\beta' < (\beta - 1) \wedge (\beta/2)$ for $\xi \in \mathfrak{M}_0$ such that $\sharp\{x \in \xi : \xi([x - |x|^{\beta'}, x + |x|^{\beta'}]) \geq 2\} = \infty$, then ξ does not satisfy the condition (C.2) (ii). In order to include such initial configurations as well as those with multiple points in our study of Dyson's model with an infinite number of particles, we introduce another condition for configurations:

(C.3) there exists $\kappa \in (1/2, 1)$ and $m \in \mathbb{N}$ such that

$$m(\xi, \kappa) \equiv \max_{k \in \mathbb{Z}} \xi \left([g^\kappa(k), g^\kappa(k+1)] \right) \leq m.$$

We denote by $\mathfrak{Y}_m^\kappa$ the set of configurations ξ satisfying (C.1) and (C.3) with $\kappa \in (1/2, 1)$ and $m \in \mathbb{N}$, and put

$$\mathfrak{Y} = \bigcup_{\kappa \in (1/2, 1)} \bigcup_{m \in \mathbb{N}} \mathfrak{Y}_m^\kappa.$$

Noting that the set $\{\xi \in \mathfrak{M} : m(\xi, \kappa) \leq m\}$ is relatively compact for each $\kappa \in (1/2, 1)$ and $m \in \mathbb{N}$, we see that $\mathfrak{Y}$ is locally compact. We introduce the following topology on $\mathfrak{Y}$.

Definition 2.3 *Suppose that $\xi, \xi_n \in \mathfrak{Y}, n \in \mathbb{N}$. We say that ξ_n converges Φ -moderately to ξ , if*

$$\lim_{n \rightarrow \infty} \Phi(\xi_n, i, \cdot) = \Phi(\xi, i, \cdot) \text{ uniformly on any compact set of } \mathbb{C}. \quad (2.4)$$

It is easy to see that (2.4) is satisfied, if ξ_n converges to ξ vaguely and the following two conditions hold:

$$\lim_{L \rightarrow \infty} \sup_{n > 0} \left| \int_{[-L, L]^c} \frac{\xi_n(dx)}{x} \right| = 0, \quad (2.5)$$

$$\lim_{L \rightarrow \infty} \sup_{n > 0} \int_{[-L, L]^c} \frac{\xi_n^{(2)}(dx)}{x} = 0. \quad (2.6)$$

Note that for any $a \in \mathbb{R}$ and $z \in \mathbb{C}$

$$\lim_{n \rightarrow \infty} \Phi(\xi_n, a, z) = \Phi(\xi, a, z), \quad (2.7)$$

if ξ_n converges Φ -moderately to ξ and $a \notin \text{supp } \xi$.

Then the second theorem of the present paper is the following.

Theorem 2.4 (i) *If $\xi \in \mathfrak{Y}$, $(\mathbb{P}_\xi, \Xi(t), t \in [0, \infty))$ is well defined with a correlation kernel $\mathbb{K}^\xi$. In particular, when $\xi \in \mathfrak{Y}_0 \equiv \mathfrak{Y} \cap \mathfrak{M}_0$, $\mathbb{K}^\xi$ is given by (2.3).*

(ii) *Suppose that $\xi, \xi_n \in \mathfrak{Y}_m^\kappa, n \in \mathbb{N}$ for some $\kappa \in (1/2, 1)$ and $m \in \mathbb{N}$. If ξ_n converges Φ -moderately to ξ , then the process $(\mathbb{P}_{\xi_n}, \Xi(t), t \in [0, \infty))$ converges to the process $(\mathbb{P}_\xi, \Xi(t), t \in [0, \infty))$ weakly in the sense of finite dimensional distributions as $n \rightarrow \infty$ in the vague topology.*

In the proof of this theorem given in Section 4.4, we will give an expression (4.26) to $\mathbb{K}^\xi$, which is valid for any $\xi \in \mathfrak{Y}$. There we will use special functions such as the Hermite polynomials, $H_k, k \in \mathbb{N}_0 \equiv \mathbb{N} \cup \{0\}$, the complete symmetric functions $h_k, k \in \mathbb{N}_0$, and the Schur functions $s_{(k|\ell)}, k, \ell \in \mathbb{N}_0$.

3 Special Functions

3.1 Multivariate symmetric functions

For $n \in \mathbb{N}$, let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$ be a partition of length less than or equal to n , and $\delta = (n-1, n-2, \dots, 1, 0)$. For $\mathbf{x} = (x_1, x_2, \dots, x_n)$ consider the skew-symmetric polynomial

$$a_{\lambda+\delta}(\mathbf{x}) = \det_{1 \leq j, k \leq n} \left[x_j^{\lambda_k + n - k} \right].$$

If $\lambda = \emptyset$, it is the Vandermonde determinant, which is given by the product of difference of variables:

$$a_\delta(\mathbf{x}) = \det_{1 \leq j, k \leq n} \left[x_j^{n-k} \right] = \prod_{1 \leq j < k \leq n} (x_j - x_k).$$

The *Schur function* of the variables $\mathbf{x} = (x_1, x_2, \dots, x_n)$ corresponding to the partition of length $\leq n$ is then defined by

$$s_\lambda(\mathbf{x}) = \frac{a_{\lambda+\delta}(\mathbf{x})}{a_\delta(\mathbf{x})},$$

which is a symmetric polynomial of $\mathbf{x}$ [13].

In the present paper, the following two special cases are considered:

(i) When $\lambda = (r)$, $s_\lambda(\mathbf{x})$ is denoted by $h_r(\mathbf{x})$ and called the r -th *complete symmetric function*, which is the sum of all monomials of total degree r in the variables $\mathbf{x} = (x_1, x_2, \dots, x_n)$. The generating function for h_r is

$$H(\mathbf{x}, z) = \sum_{r \in \mathbb{N}_0} h_r(\mathbf{x}) z^r = \prod_{j=1}^n \frac{1}{1 - x_j z} \quad \text{for } \max_{1 \leq j \leq n} |x_j z| < 1.$$

(ii) When $\lambda = (k+1, 1^\ell)$, $k + \ell + 1 \leq n$, we use Frobenius' notation $(k|\ell)$ for the partition, and consider the Schur function $s_{(k|\ell)}$. Note that the sum of coefficients of the polynomial $s_{(k|\ell)}(\mathbf{x})$ equals

$$s_{(k|\ell)}(1, \dots, 1) = \binom{k+\ell}{\ell} \binom{n}{k+\ell+1}. \quad (3.1)$$

Next we consider an infinite sequence of variables: $\mathbf{x} = (x_j)_{j \in \mathbb{N}}$. If $\sum_{j \in \mathbb{N}} x_j < \infty$, and z is a variables such that $\sup_{j \in \mathbb{N}} |x_j z| < 1$, then $|H(\mathbf{x}, z)| < \infty$. Moreover, if $\sum_{j \in \mathbb{N}} x_j^2 < \infty$ in addition to the above conditions, we can show

$$\left| \frac{d^k}{dz^k} \prod_{j \in \mathbb{N}} \frac{1}{1 - x_j z} \right|_{z=0} \leq \frac{d^k}{dz^k} \exp \left\{ \left| \sum_{j \in \mathbb{N}} x_j \right| z + \sum_{j \in \mathbb{N}} \frac{x_j^2}{1 - |x_j z|} z^2 \right\} \Big|_{z=0}, \quad k \in \mathbb{N},$$

by simple calculation. It implies

$$\sum_{r \in \mathbb{N}_0} |h_r(\mathbf{x})| z^r \leq \exp \left\{ \left| \sum_{j \in \mathbb{N}} x_j \right| z + \sum_{j \in \mathbb{N}} \frac{x_j^2}{1 - |x_j z|} z^2 \right\}, \quad (3.2)$$

and thus the formula

$$\sum_{r \in \mathbb{N}_0} h_r(\mathbf{x}) z^r = \prod_{j \in \mathbb{N}} \frac{1}{1 - x_j z} \quad (3.3)$$

is valid for the infinite sequence of variables $\mathbf{x} = (x_j)_{j \in \mathbb{N}}$. Assume that there exist $x_0 \in \mathbb{R}$ and $\varepsilon > 0$ such that $\xi([x_0 - \varepsilon, x_0 + \varepsilon]) = 0$. We see that for fixed $z \in \mathbb{C}$

$$\begin{aligned}\Phi(\xi, x, z) &= \prod_{u \in \xi - \delta_x} \left(1 - \frac{z - x}{u - x}\right) \\ &= \prod_{u \in \xi} \left(1 - \frac{z - x_0}{u - x_0}\right) \prod_{u \in \xi - \delta_x} \frac{1}{1 - (x - x_0)/(u - x_0)} \\ &= \Phi(\xi, x_0, z) \sum_{r \in \mathbb{N}_0} h_r \left(\left(\frac{1}{u - x_0} \right)_{u \in \xi} \right) (x - x_0)^r,\end{aligned}$$

where (3.3) has been used. Then $\Phi(\xi, x, z)$ is a smooth function of x on $[x_0 - \varepsilon, x_0 + \varepsilon]$.

3.2 Multiple Hermite polynomials

For any $\xi \in \mathfrak{M}$ with $\xi(\mathbb{R}) < \infty$, the *multiple Hermite polynomial of type II*, P_ξ is defined as the monic polynomial of degree $\xi(\mathbb{R})$ that satisfies for any $x \in \text{supp } \xi$

$$\int_{\mathbb{R}} dy P_\xi(y) y^j e^{-(y-x)^2/2} = 0, \quad j = 0, \dots, \xi(\{x\}) - 1. \quad (3.4)$$

The *multiple Hermite polynomials of type I* consist of a set of polynomials

$$\left\{ A_\xi(\cdot, x) : x \in \text{supp } \xi, \quad \deg A_\xi(\cdot, x) = \xi(\{x\}) - 1 \right\} \quad (3.5)$$

such that the function

$$Q_\xi(y) = \sum_{x \in \text{supp } \xi} A_\xi(y, x) e^{-(y-x)^2/2} \quad (3.6)$$

satisfies

$$\int_{\mathbb{R}} dy Q_\xi(y) y^j = \begin{cases} 0, & j = 0, \dots, \xi(\mathbb{R}) - 2 \\ 1, & j = \xi(\mathbb{R}) - 1. \end{cases} \quad (3.7)$$

The polynomials $\{A_\xi(\cdot, x)\}$ are uniquely determined by the degree requirements (3.5) and the orthogonality relations (3.7) [8]. The multiple Hermite polynomial of type II, P_ξ and the function Q_ξ defined by (3.6) have the following integration representations [2],

$$\begin{aligned}P_\xi(y) &= \int_{\mathbb{R}} dy' \frac{e^{-(y'+iy)^2/2}}{\sqrt{2\pi}} \prod_{x \in \xi} (iy' - x), \\ Q_\xi(y) &= \frac{1}{2\pi i} \oint_{\Gamma(\xi)} dz \frac{e^{-(z-y)^2/2}}{\sqrt{2\pi}} \frac{1}{\prod_{x \in \xi} (z - x)}.\end{aligned} \quad (3.8)$$

Now we fix $\xi^N \in \mathfrak{M}$ with $\xi^N(\mathbb{R}) = N \in \mathbb{N}$. We write $\xi^N(\cdot) = \sum_{j=1}^N \delta_{x_j}(\cdot)$ with a labeled configuration $\mathbf{x} = (x_j)_{j=1}^N$ such that $x_1 \leq x_2 \leq \cdots \leq x_N$. Then we define

$$\xi_0^N(\cdot) \equiv 0 \quad \text{and} \quad \xi_j^N(\cdot) = \sum_{k=1}^j \delta_{x_k}(\cdot), \quad 1 \leq j \leq N.$$

By definition $\xi_j^N(\mathbb{R}) = j, 0 \leq j \leq N$ and $\xi_j^N(\{x\}) \leq \xi_{j+1}^N(\{x\}), \forall x \in \mathbb{R}, 0 \leq j \leq N-1$. We define

$$H_j^{(-)}(y; \xi^N) = P_{\xi_j^N}(y), \quad H_j^{(+)}(y; \xi^N) = Q_{\xi_{j+1}^N}(y), \quad 0 \leq j \leq N-1. \quad (3.9)$$

By the orthogonality relations (3.4), (3.7) and the above definitions, we can prove the *biorthonormality* [2]

$$\int_{\mathbb{R}} dy H_j^{(-)}(y; \xi^N) H_k^{(+)}(y; \xi^N) = \delta_{jk}, \quad 0 \leq j, k \leq N-1. \quad (3.10)$$

For $N \in \mathbb{N}$, let $\mathbb{W}_N = \{\mathbf{x} \in \mathbb{R}^N : x_1 < x_2 < \cdots < x_N\}$, the Weyl chamber of type A_{N-1} .

Lemma 3.1 *Let $\mathbf{y} = (y_j)_{j=1}^N \in \mathbb{W}_N$. For any $\xi^N(\cdot) = \sum_{j=1}^N \delta_{x_j}(\cdot) \in \mathfrak{M}$ with a labeled configuration $\mathbf{x} = (x_j)_{j=1}^N$ such that $x_1 \leq x_2 \leq \cdots \leq x_N$,*

$$\frac{1}{a_\delta(\mathbf{x})} \det_{1 \leq j, k \leq N} \left[e^{-(y_k - x_j)^2/2} \right] = (-1)^{N(N-1)/2} (2\pi)^{N/2} \det_{1 \leq j, k \leq N} \left[H_{j-1}^{(+)}(y_k; \xi^N) \right]. \quad (3.11)$$

Here when some of the x_j 's coincide, we interpret the LHS using l'Hôpital's rule.

Proof. First we assume $\xi^N \in \mathfrak{M}_0$. Since $a_\delta(\mathbf{x}) = (-1)^{N(N-1)/2} \prod_{j=2}^N \prod_{m=1}^{j-1} (x_j - x_m)$, by the multilinearity of determinant

$$\begin{aligned} & \frac{1}{a_\delta(\mathbf{x})} \det_{1 \leq j, k \leq N} \left[e^{-(y_k - x_j)^2/2} \right] \\ &= (-1)^{N(N-1)/2} (2\pi)^{N/2} \det_{1 \leq j, k \leq N} \left[\frac{e^{-(y_k - x_j)^2/2}}{\sqrt{2\pi}} \frac{1}{\prod_{m=1}^{j-1} (x_j - x_m)} \right] \\ &= (-1)^{N(N-1)/2} (2\pi)^{N/2} \det_{1 \leq j, k \leq N} \left[\sum_{\ell=1}^j \frac{e^{-(y_k - x_\ell)^2/2}}{\sqrt{2\pi}} \frac{1}{\prod_{1 \leq m \leq j, m \neq \ell} (x_\ell - x_m)} \right]. \end{aligned}$$

By definition (3.9) with (3.8), if $\xi^N \in \mathfrak{M}_0, \xi^N(\mathbb{R}) = N$,

$$\begin{aligned} H_{j-1}^{(+)}(y_k; \xi^N) &= \frac{1}{2\pi i} \oint_{\Gamma(\xi_j^N)} dz \frac{e^{-(y_k - z)^2/2}}{\sqrt{2\pi}} \frac{1}{\prod_{x \in \xi_j^N} (z - x)} \\ &= \frac{1}{2\pi i} \oint_{\Gamma(\xi_j^N)} dz \frac{e^{-(y_k - z)^2/2}}{\sqrt{2\pi}} \frac{1}{\prod_{\ell=1}^j (z - x_\ell)} \\ &= \sum_{\ell=1}^j \frac{e^{-(y_k - x_\ell)^2/2}}{\sqrt{2\pi}} \frac{1}{\prod_{1 \leq m \leq j, m \neq \ell} (x_\ell - x_m)}, \quad 1 \leq j \leq N. \end{aligned} \quad (3.12)$$

Then (3.11) is proved for $\xi^N \in \mathfrak{M}_0$. When some of the x_j 's coincide, the LHS of (3.11) is interpreted using l'Hôpital's rule and in the RHS of (3.11) $H_{j-1}^{(+)}(y_k; \xi^N)$ should be given by (3.12). Then (3.11) is valid for any $\xi^N \in \mathfrak{M}, \xi^N(\mathbb{R}) = N$. ■

Lemma 3.2 *Let $N \in \mathbb{N}, \xi^N \in \mathfrak{M}$ with $\xi^N(\mathbb{R}) = N$. For $0 \leq s \leq t, x, y \in \mathbb{R}, 0 \leq j \leq N-1$,*

$$\int_{\mathbb{R}} dy H_j^{(-)} \left(\frac{y}{\sqrt{t}}; \frac{1}{\sqrt{t}} \circ \xi^N \right) p(t-s, y|x) = \left(\frac{s}{t} \right)^{j/2} H_j^{(-)} \left(\frac{x}{\sqrt{s}}; \frac{1}{\sqrt{s}} \circ \xi^N \right), \quad (3.13)$$

$$\int_{\mathbb{R}} dx p(t-s, y|x) H_j^{(+)} \left(\frac{x}{\sqrt{s}}; \frac{1}{\sqrt{s}} \circ \xi^N \right) = \left(\frac{s}{t} \right)^{(j+1)/2} H_j^{(+)} \left(\frac{y}{\sqrt{t}}; \frac{1}{\sqrt{t}} \circ \xi^N \right), \quad (3.14)$$

where p is the heat kernel (1.4).

Proof. Consider the integral

$$\begin{aligned} & \int_{\mathbb{R}} dy H_j^{(-)} \left(\frac{y}{\sqrt{t}}; \frac{1}{\sqrt{t}} \circ \xi^N \right) p(t-s, y|x) \\ &= \frac{1}{\sqrt{2\pi(t-s)}} \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} dy' \prod_{x \in \xi_j^N} \left(iy' - \frac{x}{\sqrt{t}} \right) \int_{\mathbb{R}} dy e^{-(y-x)^2/\{2(t-s)\} - (y' + iy/\sqrt{t})^2/2} \\ &= \sqrt{\frac{t}{s}} \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} dy' \prod_{x \in \xi_j^N} \left(iy' - \frac{x}{\sqrt{t}} \right) e^{-t(y' + ix/\sqrt{t})^2/(2s)}. \end{aligned}$$

Change the integral variable $y' \rightarrow y' \sqrt{t/s}$ to obtain the equality (3.13). Similar calculation gives (3.14). ■

When $\xi^N(\cdot) = N\delta_0(\cdot)$,

$$\begin{aligned} H_j^{(-)}(y; N\delta_0) &= 2^{-j/2} H_j(y/\sqrt{2}), \\ H_j^{(+)}(y; N\delta_0) &= \frac{2^{-j/2}}{j! \sqrt{2\pi}} H_j(y/\sqrt{2}) e^{-y^2/2}, \quad 0 \leq j \leq N-1, \end{aligned}$$

where $H_j(x)$ is the Hermite polynomial of degree j ,

$$\begin{aligned} H_j(x) &= j! \sum_{k=0}^{\lfloor j/2 \rfloor} (-1)^k \frac{(2x)^{j-2k}}{k!(j-2k)!} \\ &= 2^{j/2} \int_{\mathbb{R}} dy \frac{e^{-y^2/2}}{\sqrt{2\pi}} (iy + \sqrt{2}x)^j \\ &= \frac{j!}{2\pi i} \oint_{\Gamma(\delta_0)} dz \frac{e^{2zx-z^2}}{z^{j+1}}. \end{aligned} \quad (3.15)$$

The last expression (3.15) implies that the generating function of the Hermite polynomials is given by

$$e^{2zx-z^2} = \sum_{j \in \mathbb{N}_0} \frac{z^j}{j!} H_j(x). \quad (3.16)$$

4 Proofs of Results

4.1 Proof of Proposition 2.1

For $\mathbf{x}, \mathbf{y} \in \mathbb{W}_N$ and $t > 0$, consider the *Karlin-McGregor determinant* of the heat kernel (1.4) [9]

$$f_N(t, \mathbf{y} | \mathbf{x}) = \det_{1 \leq j, k \leq N} [p(t, y_j | x_k)].$$

If $\xi^N \in \mathfrak{M}_0$ with $\xi^N(\mathbb{R}) = N \in \mathbb{N}$, ξ^N can be identified with a set $\mathbf{x} \in \mathbb{W}_N$. For any $M \geq 1$ and any time sequence $0 < t_1 < \dots < t_M < \infty$, the *multitime probability density* of Dyson's model is given by [6, 10]

$$\begin{aligned} p^{\xi^N}(t_1, \xi^{(1)}; \dots; t_M, \xi^{(M)}) \\ = a_\delta(\mathbf{x}^{(M)}) \prod_{m=1}^{M-1} f_N(t_{m+1} - t_m, \mathbf{x}^{(m+1)} | \mathbf{x}^{(m)}) f_N(t_1, \mathbf{x}^{(1)} | \mathbf{x}) \frac{1}{a_\delta(\mathbf{x})}, \end{aligned}$$

where $\xi^{(m)}(\cdot) = \sum_{j=1}^N \delta_{x_j^{(m)}}(\cdot)$, $1 \leq m \leq M$.

Define

$$\begin{aligned} \phi_j^{(-)}(t, x; \xi^N) &\equiv t^{j/2} H_j^{(-)}\left(\frac{x}{\sqrt{t}}; \frac{1}{\sqrt{t}} \circ \xi^N\right), \\ \phi_j^{(+)}(t, x; \xi^N) &\equiv t^{-(j+1)/2} H_j^{(+)}\left(\frac{x}{\sqrt{t}}; \frac{1}{\sqrt{t}} \circ \xi^N\right), \end{aligned}$$

$0 \leq j \leq N-1, t > 0, x \in \mathbb{R}$. From the biorthonormality (3.10) of the multiple Hermite polynomials and Lemma 3.2, the following relations are derived.

Lemma 4.1 *For $\xi^N \in \mathfrak{M}$ with $\xi^N(\mathbb{R}) = N \in \mathbb{N}$, $0 \leq t_1 \leq t_2$,*

$$\begin{aligned} \int_{\mathbb{R}} dx_2 \phi_j^{(-)}(t_2, x_2; \xi^N) p(t_2 - t_1, x_2 | x_1) &= \phi_j^{(-)}(t_1, x_1; \xi^N), \quad 0 \leq j \leq N-1, \\ \int_{\mathbb{R}} dx_1 p(t_2 - t_1, x_2 | x_1) \phi_j^{(+)}(t_1, x_1; \xi^N) &= \phi_j^{(+)}(t_2, x_2; \xi^N), \quad 0 \leq j \leq N-1, \\ \int_{\mathbb{R}} dx_1 \int_{\mathbb{R}} dx_2 \phi_j^{(-)}(t_2, x_2; \xi^N) p(t_2 - t_1, x_2 | x_1) \phi_k^{(+)}(t_1, x_1; \xi^N) &= \delta_{jk}, \\ &0 \leq j, k \leq N-1. \end{aligned}$$

Put

$$\mu^{(\pm)}(t, \mathbf{x}; \xi^N) = \det_{1 \leq j, k \leq N} \left[\phi_{j-1}^{(\pm)}(t, x_k; \xi^N) \right].$$

Since $H_j^{(-)}$ is a monic polynomial of degree j , $\mu^{(-)}(t, \mathbf{x}; \xi^N) = (-1)^{N(N-1)/2} a_\delta(\mathbf{x})$. By Lemma 3.1, $f_N(t_1, \mathbf{x}^{(1)} | \mathbf{x}) / a_\delta(\mathbf{x})$ will be replaced by $(-1)^{N(N-1)/2} \mu^{(+)}(t_1, \mathbf{x}^{(1)}; \xi^N)$ to extend the expression to the case $\xi^N \in \mathfrak{M}$. Then the multitime probability density of Dyson's model is expressed as

$$\begin{aligned} & p^{\xi^N} \left(t_1, \xi^{(1)}; \dots; t_M, \xi^{(M)} \right) \\ &= \mu^{(-)}(t_M, \mathbf{x}^{(M)}; \xi^N) \prod_{m=1}^{M-1} f_N(t_{m+1} - t_m; \mathbf{x}^{(m+1)} | \mathbf{x}^{(m)}) \mu^{(+)}(t_1, \mathbf{x}^{(1)}; \xi^N) \end{aligned} \quad (4.1)$$

for $\xi^N \in \mathfrak{M}$ with $\xi^N(\mathbb{R}) = N \in \mathbb{N}$. For $\mathbf{x} = (x_1, \dots, x_N)$ with $\xi(\cdot) = \sum_{j=1}^N \delta_{x_j}(\cdot)$ and $N' \in \{1, 2, \dots, N\}$, we put $\mathbf{x}_{N'} = (x_1, \dots, x_{N'})$. For a sequence $(N_m)_{m=1}^M$ of positive integers less than or equal to N , we obtain the $(N_1, \dots, N_M)$ -multitime correlation function by

$$\begin{aligned} & \rho_N^{\xi^N} \left(t_1, \mathbf{x}_{N_1}^{(1)}; \dots; t_M, \mathbf{x}_{N_M}^{(M)} \right) \\ &= \int_{\prod_{m=1}^M \mathbb{R}^{N-N_m}} p^{\xi^N} \left(t_1, \xi^{(1)}; \dots; t_M, \xi^{(M)} \right) \prod_{m=1}^M \frac{1}{(N - N_m)!} \prod_{j=N_m+1}^N dx_j^{(m)}. \end{aligned} \quad (4.2)$$

For $\mathbf{f} = (f_1, \dots, f_M) \in C_0(\mathbb{R})^M$, and $\boldsymbol{\theta} = (\theta_1, \dots, \theta_M) \in \mathbb{R}^M$, the generating function for multitime correlation functions is given as

$$\begin{aligned} \mathcal{G}^{\xi^N}[\chi] &= \mathbb{E}_{\xi^N} \left[\exp \left\{ \sum_{m=1}^M \theta_m \sum_{j_m=1}^N f_m(X_{j_m}(t_m)) \right\} \right] \\ &= \sum_{N_1=0}^N \dots \sum_{N_M=0}^N \prod_{m=1}^M \frac{1}{N_m!} \int_{\mathbb{R}^{N_1}} \prod_{j=1}^{N_1} dx_j^{(1)} \dots \int_{\mathbb{R}^{N_M}} \prod_{j=1}^{N_M} dx_j^{(M)} \\ &\quad \times \prod_{m=1}^M \prod_{j=1}^{N_m} \chi_m(x_j^{(m)}) \rho^{\xi^N} \left(t_1, \mathbf{x}_{N_1}^{(1)}; \dots; t_M, \mathbf{x}_{N_M}^{(M)} \right), \end{aligned}$$

where

$$\chi_m(x) = e^{\theta_m f_m(x)} - 1, \quad 1 \leq m \leq M.$$

By the argument given in Section 4.2 in [10], the expression (4.1) with Lemma 4.1 leads to the Fredholm determinantal expression for the generating function,

$$\mathcal{G}^{\xi^N}[\chi] = \text{Det} \left[\delta_{mn} \delta(x - y) + \tilde{S}^{m,n}(x, y; \xi^N) \chi_n(y) \right],$$

where

$$\tilde{S}^{m,n}(x, y; \xi^N) = S^{m,n}(x, y; \xi^N) - \mathbf{1}(m > n) p(t_m - t_n, x | y)$$

with

$$\begin{aligned}
S^{m,n}(x, y; \xi^N) &= \sum_{j=0}^{N-1} \phi_j^{(+)}(t_m, x; \xi^N) \phi_j^{(-)}(t_n, y; \xi^N) \\
&= \frac{1}{\sqrt{t_m}} \sum_{j=0}^{N-1} \left(\frac{t_n}{t_m} \right)^{j/2} H_j^{(+)} \left(\frac{x}{\sqrt{t_m}}; \frac{1}{\sqrt{t_m}} \circ \xi^N \right) H_j^{(-)} \left(\frac{y}{\sqrt{t_n}}; \frac{1}{\sqrt{t_n}} \circ \xi^N \right).
\end{aligned}$$

Here the Fredholm determinant is expanded as

$$\begin{aligned}
&\text{Det} \left[\delta_{mn} \delta(x - y) + \tilde{S}^{m,n}(x, y; \xi^N) \chi_n(y) \right] \\
&= \sum_{N_1=0}^N \cdots \sum_{N_M=0}^N \prod_{m=1}^M \frac{1}{N_m!} \int_{\mathbb{R}^{N_1}} \prod_{j=1}^{N_1} dx_j^{(1)} \cdots \int_{\mathbb{R}^{N_M}} \prod_{j=1}^{N_M} dx_j^{(M)} \\
&\quad \times \prod_{m=1}^M \prod_{j=1}^{N_m} \chi_m(x_j^{(m)}) \det_{\substack{1 \leq j \leq N_m, 1 \leq k \leq N_n \\ 1 \leq m, n \leq M}} \left[\tilde{S}^{m,n}(x_j^{(m)}, x_k^{(n)}; \xi^N) \right].
\end{aligned}$$

Proof of Proposition 2.1. Inserting the integral formulas for $H_j^{(\pm)}$, the kernel $S^{m,n}$ is written as

$$\begin{aligned}
S^{m,n}(x, y; \xi^N) &= \frac{1}{\sqrt{t_m}} \frac{1}{2\pi i} \oint_{\Gamma(t_m^{-1/2} \circ \xi^N)} dz \frac{e^{-(z-x/\sqrt{t_m})^2/2}}{\sqrt{2\pi}} \int_{\mathbb{R}} dy' \frac{e^{-(y'+iy/\sqrt{t_n})^2/2}}{\sqrt{2\pi}} \\
&\quad \times \sum_{k=0}^{N-1} \left(\frac{t_n}{t_m} \right)^{k/2} \frac{\prod_{\ell=1}^k (iy' - x_\ell/\sqrt{t_n})}{\prod_{\ell=1}^{k+1} (z - x_\ell/\sqrt{t_m})} \\
&= \frac{1}{2\pi i} \oint_{\Gamma(t_m^{-1/2} \circ \xi^N)} dz \frac{e^{-(z-x/\sqrt{t_m})^2/2}}{\sqrt{2\pi}} \int_{\mathbb{R}} dy' \frac{e^{-(y'+iy/\sqrt{t_n})^2/2}}{\sqrt{2\pi}} \\
&\quad \times \sum_{k=0}^{N-1} \frac{\prod_{\ell=1}^k (i\sqrt{t_n}y' - x_\ell)}{\prod_{\ell=1}^{k+1} (\sqrt{t_m}z - x_\ell)}.
\end{aligned}$$

For $z_1, z_2 \in \mathbb{C}$ with $z_1 \notin \{x_1, \dots, x_N\}$, the following identity holds,

$$\begin{aligned}
&\sum_{k=0}^{N-1} \frac{\prod_{\ell=1}^k (z_2 - x_\ell)}{\prod_{\ell=1}^{k+1} (z_1 - x_\ell)} \\
&= \frac{1}{z_1 - x_1} + \frac{z_2 - x_1}{(z_1 - x_1)(z_1 - x_2)} + \cdots + \frac{(z_2 - x_1)(z_2 - x_2) \cdots (z_2 - x_{N-1})}{(z_1 - x_1)(z_1 - x_2) \cdots (z_1 - x_N)} \\
&= \left(\prod_{\ell=1}^N \frac{z_2 - x_\ell}{z_1 - x_\ell} - 1 \right) \frac{1}{z_2 - z_1}.
\end{aligned}$$

By this identity, we have

$$\begin{aligned} S^{m,n}(x, y; \xi^N) &= \frac{1}{2\pi i} \oint_{\Gamma(t_m^{-1/2} \circ \xi^N)} dz \frac{e^{-(z-x/\sqrt{t_m})^2/2}}{\sqrt{2\pi}} \int_{\mathbb{R}} dy' \frac{e^{-(y'+iy/\sqrt{t_n})^2/2}}{\sqrt{2\pi}} \\ &\quad \times \left(\prod_{\ell=1}^N \frac{i\sqrt{t_n}y' - x_\ell}{\sqrt{t_m}z - x_\ell} - 1 \right) \frac{1}{i\sqrt{t_n}y' - \sqrt{t_m}z}. \end{aligned}$$

Note that

$$\begin{aligned} &\frac{1}{2\pi i} \oint_{\Gamma(t_m^{-1/2} \circ \xi^N)} dz \frac{e^{-(z-x/\sqrt{t_m})^2/2}}{\sqrt{2\pi}} \int_{\mathbb{R}} dy' \frac{e^{-(y'+iy/\sqrt{t_n})^2/2}}{\sqrt{2\pi}} \frac{1}{i\sqrt{t_n}y' - \sqrt{t_m}z} \\ &= \frac{1}{2\pi i} \oint_{\Gamma(t_m^{-1/2} \circ \xi^N)} dz \frac{e^{-(z-x/\sqrt{t_m})^2/2}}{\sqrt{2\pi}} \int_{\mathbb{R}} dy' \frac{e^{-(y'+iy/\sqrt{t_n})^2/2}}{\sqrt{2\pi}} \frac{1}{i\sqrt{t_n}y'} \sum_{j \in \mathbb{N}_0} \left(\sqrt{\frac{t_m}{t_n}} \frac{z}{iy'} \right)^j \\ &= 0. \end{aligned}$$

By changing the integral variables appropriately, we find that $\tilde{S}^{m,n}(x, y; \xi^N)$ is equal to (2.1) with $s = t_m, t = t_n$. This completes the proof. ■

4.2 Proof of Theorem 2.2

In this subsection we give a proof of Theorem 2.2. First we prove some lemmas.

Lemma 4.2 *If $M_\alpha(\xi) < \infty$ for some $\alpha \in (1, 2)$, then*

$$\alpha \sum_{L \in \mathbb{N}} \frac{M_1(\xi, L)^{\alpha/(\alpha-1)}}{L(L+1)^\alpha} \leq M_\alpha(\xi)^{\alpha^2/(\alpha-1)}.$$

Proof. By Hölder's inequality we have

$$M_1(\xi, L) = \int_{0 < |x| \leq L} \frac{\xi(dx)}{|x|} \leq M_\alpha(\xi) \xi([-L, L] \setminus \{0\})^{(\alpha-1)/\alpha}.$$

On the other hand

$$\begin{aligned} M_\alpha(\xi)^\alpha &= \sum_{L \in \mathbb{N}} \int_{L-1 < |x| \leq L} \frac{\xi(dx)}{|x|^\alpha} \\ &\geq \sum_{L \in \mathbb{N}} L^{-\alpha} \left\{ \xi([-L, L] \setminus \{0\}) - \xi([-L+1, L-1] \setminus \{0\}) \right\} \\ &= \sum_{L \in \mathbb{N}} \left\{ L^{-\alpha} - (L+1)^{-\alpha} \right\} \xi([-L, L] \setminus \{0\}) \\ &\geq \alpha \sum_{L \in \mathbb{N}} \frac{\xi([-L, L] \setminus \{0\})}{L(L+1)^\alpha}. \end{aligned}$$

From the above inequalities we have

$$M_\alpha(\xi)^\alpha \geq \alpha \sum_{L \in \mathbb{N}} \frac{1}{L(L+1)^\alpha} \left(\frac{M_1(\xi, L)}{M_\alpha(\xi)} \right)^{\alpha/(\alpha-1)}.$$

Lemma 4.2 is derived from this inequality, since $\alpha + \alpha/(\alpha-1) = \alpha^2/(\alpha-1)$. ■

Lemma 4.3 *Let $\alpha \in (1, 2)$ and $\delta > \alpha - 1$. Suppose that $M_\alpha(\xi) < \infty$ and put $L_0 = L_0(\alpha, \delta, \xi) = (2M_\alpha(\xi))^{\alpha/(\delta-\alpha+1)}$. Then*

$$M_1(\xi, L) \leq L^\delta, \quad L \geq L_0.$$

Proof. Suppose that $L_1 \in \mathbb{N}$ satisfies $M_1(\xi, L_1) > L_1^\delta$. Then

$$\begin{aligned} \alpha \sum_{L \in \mathbb{N}} \frac{M_1(\xi, L)^{\alpha/(\alpha-1)}}{L(L+1)^\alpha} &> \alpha \sum_{L=L_1}^{\infty} \frac{L_1^{\alpha\delta/(\alpha-1)}}{L(L+1)^\alpha} \\ &> \alpha L_1^{\alpha\delta/(\alpha-1)} \int_{L_1+1}^{\infty} dy y^{-(\alpha+1)} \\ &= L_1^{\alpha\delta/(\alpha-1)} (L_1+1)^{-\alpha} = \left(\frac{L_1}{L_1+1} \right)^\alpha L_1^{\alpha(\delta-\alpha+1)/(\alpha-1)}. \end{aligned}$$

From Lemma 4.2 we have

$$\left(\frac{L_1}{L_1+1} \right)^\alpha L_1^{\alpha(\delta-\alpha+1)/(\alpha-1)} \leq M_\alpha(\xi)^{\alpha^2/(\alpha-1)}.$$

Hence

$$L_1 < \left(\frac{L_1+1}{L_1} \right)^{(\alpha-1)/(\delta-\alpha+1)} M_\alpha(\xi)^{\alpha/(\delta-\alpha+1)} < (2M_\alpha(\xi))^{\alpha/(\delta-\alpha+1)}.$$

This completes the proof. ■

The following lemma will play an important role in the proof of Theorem 2.2.

Lemma 4.4 *For any $\xi \in \mathfrak{X}_0$, there exist $C_3 = C_3(\alpha, \beta, C_0, C_1, C_2) > 0$ and $\theta \in (\alpha \vee (2-\beta), 2)$ such that*

$$|\Phi(\xi, a, iy)| \leq \exp \left[C_3 \left\{ (|y|^\theta \vee 1) + (|a|^\theta \vee 1) \right\} \right] \quad \forall y \in \mathbb{R}, \quad \forall a \in \text{supp } \xi.$$

Proof. First we estimate the entire function $\Phi(\xi, a, z), z \in \mathbb{C}$, in the case that $a = 0 \in \text{supp } \xi$. In case $2|z| < |x|$, by using the expansion

$$\log \left(1 + \frac{z}{x} \right) = \sum_{k \in \mathbb{N}} \frac{(-1)^{k-1}}{k} \left(\frac{z}{x} \right)^k,$$

we have

$$\begin{aligned} \int_{2|z| < |x|} \xi(dx) \log \left(1 + \frac{z}{x} \right) &= \int_{2|z| < |x|} \xi(dx) \sum_{k \in \mathbb{N}} \frac{(-1)^{k-1}}{k} \left(\frac{z}{x} \right)^k \\ &= \int_{2|z| < |x|} \xi(dx) \frac{z}{x} + \int_{2|z| < |x|} \xi(dx) \left(\frac{z}{x} \right)^2 \sum_{k=2}^{\infty} \frac{(-1)^{k-1}}{k} \left(\frac{z}{x} \right)^{k-2}. \end{aligned}$$

Since

$$\left| \int_{2|z| < |x|} \xi(dx) \frac{z}{x} \right| \leq |M(\xi)| |z| + M_1(\xi, 2|z|) |z|,$$

and

$$\begin{aligned} &\left| \int_{2|z| < |x|} \xi(dx) \left(\frac{z}{x} \right)^2 \sum_{k=2}^{\infty} \frac{(-1)^{k-1}}{k} \left(\frac{z}{x} \right)^{k-2} \right| \\ &\leq \int_{2|z| < |x|} \xi(dx) \frac{|z|^2}{|x|^2} \frac{1}{2} \sum_{k=2}^{\infty} 2^{2-k} = \int_{2|z| < |x|} \xi(dx) \frac{|z|^2}{|x|^2} \\ &\leq M_\alpha(\xi)^\alpha |z|^\alpha, \end{aligned}$$

we have

$$\prod_{x \in \xi} \left\{ 1 + \mathbf{1}(|x| > 2|z|) \frac{z}{x} \right\} \leq \exp \left\{ |M(\xi)| |z| + M_1(\xi, 2|z|) |z| + M_\alpha(\xi)^\alpha |z|^\alpha \right\}. \quad (4.3)$$

On the other hand we have

$$\begin{aligned} &\prod_{x \in \xi} \left\{ 1 + \mathbf{1}(0 < |x| \leq 2|z|) \frac{z}{x} \right\} \\ &\leq \exp \left\{ \int_{[-2|z|, 2|z|] \setminus \{0\}} \frac{\xi(dx)}{|x|} |z| \right\} = \exp \left\{ M_1(\xi, 2|z|) |z| \right\}. \end{aligned} \quad (4.4)$$

Combining the above two inequalities (4.3) and (4.4), we obtain

$$\prod_{x \in \xi \cap \{0\}^c} \left(1 + \frac{z}{x} \right) \leq \exp \left\{ |M(\xi)| |z| + 2M_1(\xi, 2|z|) |z| + M_\alpha(\xi)^\alpha |z|^\alpha \right\}.$$

By the conditions **(C.1)**, **(C.2)**(i) and Lemma 4.3, we have

$$|M(\xi)| |z| + 2M_1(\xi, 2|z|) |z| + M_\alpha(\xi)^\alpha |z|^\alpha \leq C_0 |z| + 4|z|^{1+\delta} + C_1 |z|^\alpha$$

for $|z| \leq L_0(\alpha, \delta, C_1)$ with $\delta > \alpha - 1$. Hence, if we can take $\theta > \alpha$, then

$$|\Phi(\xi, 0, z)| \leq \exp \left[C(|z|^\theta \vee 1) \right] \quad (4.5)$$

with a positive constant C , which depends on only α, β, C_0 and C_1 .

Next we estimate the case that $a \neq 0$ by using the following equations:

$$\Phi(\xi, a, z) = \Phi(\xi, 0, z)\Phi(\xi \cap \{0\}^c, a, 0) \left(\frac{z}{a}\right)^{\xi(\{0\})} \frac{a}{a-z}$$

and

$$\Phi(\xi \cap \{0\}^c, a, 0) = \Phi(\xi \cap \{-a\}^c, 0, -a)\Phi(\xi^{(2)} \cap \{0\}^c, a^2, 0)2^{1-\xi(\{-a\})},$$

where $a \in \text{supp } \xi$. From the above estimate (4.5) we have

$$|\Phi(\xi \cap \{-a\}^c, 0, -a)| \leq \exp \left[C(|a|^\theta \vee 1) \right].$$

Since $|(iy/a)^{\xi(\{0\})}a/(a-iy)| \leq 1$, it is enough to show the estimate

$$\Phi(\xi^{(2)} \cap \{0\}^c, a^2, 0) \leq \exp \left\{ 3C_2|a|^{2-\beta} \right\}$$

for proving this lemma. We have

$$\begin{aligned} & \left| \int_{2a^2 < |x-a^2|} \xi^{(2)}(dx) \log \left(1 + \frac{a^2}{x-a^2} \right) \right| \\ &= \left| \int_{2a^2 < |x-a^2|} \xi^{(2)}(dx) \frac{a^2}{x-a^2} \sum_{k \in \mathbb{N}} \frac{(-1)^k}{k} \left(\frac{a^2}{x-a^2} \right)^{k-1} \right| \\ &\leq 2 \int_{2a^2 < |x-a^2|} \xi^{(2)}(dx) \left| \frac{a^2}{x-a^2} \right| \leq 2M_1(\tau_{-a^2}\xi^{(2)})a^2. \end{aligned}$$

On the other hand we see

$$\begin{aligned} & \prod_{x \in \xi^{(2)}} \left\{ 1 + \mathbf{1}(0 < |x-a^2| < 2a^2) \frac{a^2}{x-a^2} \right\} \\ &\leq \exp \left\{ \int_{[-2a^2, 2a^2] \setminus \{0\}} \frac{(\tau_{-a^2}\xi^{(2)})(dx)}{|x|} a^2 \right\} = \exp \left\{ M_1(\tau_{-a^2}\xi^{(2)}, 2a^2)a^2 \right\}. \end{aligned}$$

Then

$$\begin{aligned} \Phi(\xi^{(2)} \cap \{0\}^c, a^2, 0) &= \prod_{x \in \xi^{(2)} \cap \{0, a^2\}^c} \left(1 + \frac{a^2}{x-a^2} \right) \\ &\leq \exp \left\{ 3M_1(\tau_{-a^2}\xi^{(2)})a^2 \right\} = \exp \left\{ 3C_2|a|^{2-\beta} \right\}. \end{aligned}$$

This completes the proof. ■

Proof of Theorem 2.2. Note that $\xi \cap [-L, L]$, $L > 0$ and ξ satisfy (C.1) and (C.2) with the same constants C_0, C_1, C_2 and indices α, β . By virtue of Lemma 4.4 we see that there exists $C_3 > 0$ such that

$$|\Phi(\xi \cap [-L, L], a, iy)| \leq \exp \left[C_3 \left\{ |y|^\theta + (|a|^\theta \vee 1) \right\} \right],$$

$\forall L > 0, \forall a \in \text{supp } \xi, \forall y \in \mathbb{R}$. Since for any $y \in \mathbb{R}$

$$\Phi(\xi \cap [-L, L], a, iy) \rightarrow \Phi(\xi, a, iy), \quad L \rightarrow \infty,$$

we can apply Lebesgue's convergence theorem to (2.2) and obtain

$$\lim_{L \rightarrow \infty} \mathbb{K}^{\xi \cap [-L, L]}(s, x; t, y) = \mathbb{K}^\xi(s, x; t, y).$$

Since for any $(s, t) \in (0, \infty)^2$ and any compact interval $I \subset \mathbb{R}$

$$\sup_{x, y \in I} \left| \mathbb{K}^{\xi \cap [-L, L]}(s, x; t, y) \right| < \infty,$$

we can obtain the convergence of generating functions for multitime correlation functions; $\mathcal{G}^{\xi \cap [-L, L]}[\chi] \rightarrow \mathcal{G}^\xi[\chi]$ as $L \rightarrow \infty$. It implies $\mathbb{P}_{\xi \cap [-L, L]} \rightarrow \mathbb{P}_\xi$ as $L \rightarrow \infty$ in the sense of finite dimensional distributions and the proof is completed. ■

4.3 Proofs of (1.5) and (1.7)

Proof of (1.5). Since $\xi^\mathbb{Z} = \eta^1 \in \mathfrak{X}_0$, we can start from the expression of the correlation kernel (2.3) in Theorem 2.2. For $\ell \in \mathbb{Z}, z \in \mathbb{C}$

$$\begin{aligned} \Phi(\xi^\mathbb{Z}, \ell, z) &= \prod_{j \in \mathbb{Z}, j \neq \ell} \left(1 - \frac{z - \ell}{j - \ell} \right) \\ &= \frac{\sin\{\pi(z - \ell)\}}{\pi(z - \ell)} \\ &= \frac{1}{2\pi} \int_{|k| \leq \pi} dk e^{ik(z - \ell)}, \end{aligned}$$

since $\prod_{n \in \mathbb{N}} (1 - x^2/n^2) = \sin(\pi x)/(\pi x)$. Then

$$\mathbb{K}^{\xi^\mathbb{Z}}(s, x; t, y) + \mathbf{1}(s > t)p(s - t, x|y) = \sum_{\ell \in \mathbb{Z}} p(s, x|\ell)I(t, y, \ell), \quad (4.6)$$

where

$$\begin{aligned} I(t, y, \ell) &= \int_{\mathbb{R}} dy' p(t, -iy|y') \frac{1}{2\pi} \int_{|k| \leq \pi} dk e^{ik(iy' - \ell)} \\ &= \frac{1}{2\pi} \int_{|k| \leq \pi} dk e^{k^2 t/2 + ik(y - \ell)}. \end{aligned}$$

By definition (1.6) of ϑ_3 , we can rewrite (4.6) as

$$\begin{aligned} &\frac{1}{2\pi} \int_{|k| \leq \pi} dk e^{k^2(t-s)/2 + ik(y-x)} \\ &\times \vartheta_3 \left(\frac{1}{2\pi i s} (x - iks), -\frac{1}{2\pi i s} \right) e^{-\pi i (x - iks)^2 / (2\pi i s)} \sqrt{\frac{i}{2\pi i s}}. \end{aligned}$$

Use the functional equation satisfied by $\vartheta_3(v, \tau)$ (see, for example, Section 10.12 in [1]),

$$\vartheta_3(v, \tau) = \vartheta_3\left(\frac{v}{\tau}, -\frac{1}{\tau}\right) e^{-\pi i v^2 / \tau} \sqrt{\frac{i}{\tau}},$$

and the integral representation of the heat kernel (1.4). Then (1.5) is obtained. ■

Proof of (1.7). By the definition (1.6) of ϑ_3 , for $s, t, u > 0$

$$\begin{aligned} & \mathbb{K}^{\xi^{\mathbb{Z}}}(u + s, x; u + t, y) - \mathbf{K}_{\sin}(t - s, y - x) \\ &= \frac{e^{-2\pi i x}}{2\pi} \int_{|k| \leq \pi} dk e^{k^2(t-s)/2 + ik(y-x) - 2\pi(u+s)(\pi+k)} \\ & \quad + \frac{e^{2\pi i x}}{2\pi} \int_{|k| \leq \pi} dk e^{k^2(t-s)/2 + ik(y-x) - 2\pi(u+s)(\pi-k)} \\ & \quad + \sum_{\ell \in \mathbb{Z} \setminus \{-1, 0, 1\}} \frac{e^{2\pi i x \ell}}{2\pi} \int_{|k| \leq \pi} dk e^{k^2(t-s)/2 + ik(y-x) - 2\pi(u+s)\ell(\ell\pi-k)}. \end{aligned}$$

Then we see for any $u > 0$

$$\begin{aligned} & \left| \mathbb{K}^{\xi^{\mathbb{Z}}}(u + s, x; u + t, y) - \mathbf{K}_{\sin}(t - s, y - x) \right| \\ & \leq \left(e^{\pi^2(t-s)/2} \vee 1 \right) \left\{ \frac{1}{\pi} \int_{|k| \leq \pi} dk e^{-2\pi(u+s)(\pi+k)} + 2 \sum_{\ell \geq 2} e^{-2\pi^2(u+s)\ell} \right\} \\ & = \left(e^{\pi^2(t-s)/2} \vee 1 \right) \left\{ \frac{1 - e^{-4\pi^2(u+s)}}{2\pi^2(u+s)} + \frac{2e^{-4\pi^2(u+s)}}{1 - e^{-2\pi^2(u+s)}} \right\} \leq \frac{C}{u}, \end{aligned}$$

where $C > 0$ depends on t and s , but does not on u . This completes the proof of (1.7). ■

Remark Since this relaxation process $(\mathbb{P}_{\xi^{\mathbb{Z}}}, \Xi(t), t \in [0, \infty))$ is determinantal with $\mathbb{K}^{\xi^{\mathbb{Z}}}$, at any intermediate time $0 < t < \infty$ the particle distribution on $\mathbb{R}$ is the determinantal point process with the spatial correlation kernel $\mathbb{K}^{\xi^{\mathbb{Z}}}(x, t; y, t), x, y \in \mathbb{R}$. It should be noted that this spatial correlation kernel is not symmetric,

$$\mathbb{K}^{\xi^{\mathbb{Z}}}(t, x; t, y) = \sum_{\ell \in \mathbb{Z}} e^{2\pi i x \ell - 2\pi^2 t \ell^2} \frac{\sin \left[\pi \{ (y - x) - 2\pi i t \ell \} \right]}{\pi \{ (y - x) - 2\pi i t \ell \}},$$

$x, y \in \mathbb{R}, 0 < t < \infty$.

4.4 Proof of Theorem 2.4

In this subsection we prove Theorem 2.4. Suppose $\xi \in \mathfrak{Y}_m^\kappa \subset \mathfrak{Y}$. For $k \in \mathbb{Z}$ we can take $\underline{b}_k$ and $\bar{b}_k$ such that $\underline{b}_{-k-1} = -\bar{b}_k$, $\bar{b}_{-k-1} = -\underline{b}_k$,

$$\begin{aligned} [\underline{b}_k, \bar{b}_k] &\subset (g^\kappa(k), g^\kappa(k+1)), \quad \bar{b}_k - \underline{b}_k \geq \frac{g^\kappa(k+1) - g^\kappa(k)}{2m(\xi, \kappa) + 1}, \\ \xi([\underline{b}_k, \bar{b}_k]) &= 0 \quad \text{and} \quad \xi(\{(\bar{b}_{k-1} + \underline{b}_k)/2\}) = 0. \end{aligned}$$

We put $I_k = [\underline{b}_k, \bar{b}_k]$, $\varepsilon_k = |I_k| = \bar{b}_k - \underline{b}_k$, $c_k = (\bar{b}_{k-1} + \underline{b}_k)/2$, and $\Delta_k = (\underline{b}_k - \bar{b}_{k-1})/2$. Note that $[\bar{b}_{-k-1}, \underline{b}_{-k}] = -[\bar{b}_{k-1}, \underline{b}_k]$, $I_{-k-1} = -I_k$, $\varepsilon_{-k-1} = \varepsilon_k$, $k \in \mathbb{N}_0$. Then we define the k -th *cluster* in the configuration ξ by

$$\mathfrak{C}_k = \xi \cap [\bar{b}_{k-1}, \underline{b}_k].$$

It is easy to see that $\sum_{k \in \mathbb{Z}} \mathfrak{C}_k = \xi$, and for each $k \in \mathbb{Z}$

$$|\mathfrak{C}_k| \equiv \mathfrak{C}_k(\mathbb{R}) = \xi([\bar{b}_{k-1}, \underline{b}_k]) \leq 2m(\xi, \kappa), \quad (4.7)$$

$$|x - y| \geq \varepsilon_{k-1} \wedge \varepsilon_k, \quad x \in \text{supp } \mathfrak{C}_k, \quad y \in \text{supp } (\xi - \mathfrak{C}_k). \quad (4.8)$$

Let $\mathbf{v}_k = (v_{k\ell})_{\ell=1}^{|\mathfrak{C}_k|}$ be the increasing sequence with $\sum_{\ell=1}^{|\mathfrak{C}_k|} \delta_{v_{k\ell}} = \mathfrak{C}_k$. See Figure 1. For $a \in \text{supp } \xi$, we denote by $\mathfrak{C}^a$ the cluster containing a . Remark that, when ξ_n converges to ξ vaguely as $n \rightarrow \infty$, we can take the k -th cluster $\mathfrak{C}_k(\xi_n)$ of ξ_n so that it converges to the k -th cluster $\mathfrak{C}_k(\xi)$ of ξ vaguely as $n \rightarrow \infty$.




Figure 1: The clusters

We introduce $\mathbb{C}$ -valued functions $\Psi_k(t, \xi, z, x)$, $k \in \mathbb{Z}$, $t \geq 0$, $\xi \in \mathfrak{Y}$, $z \in \mathbb{C}$,

$x \in \mathbb{R}$,

$$\begin{aligned} \Psi_k(t, \xi, z, x) &= \Phi(\xi - \mathfrak{C}_k, c_k, z) \sum_{\ell=1}^{|\mathfrak{C}_k|} (z - c_k)^{\ell-1} (-1)^{|\mathfrak{C}_k|-\ell-1} \\ &\times \left\{ \Theta_{k,\ell}(t, \xi, x) + \sum_{q=|\mathfrak{C}_k|}^{\infty} \Theta_{k,q}(t, \xi, x) s_{(q-|\mathfrak{C}_k|, |\mathfrak{C}_k|-\ell-1)}(\mathbf{v}_k - c_k) \right\}, \end{aligned} \quad (4.9)$$

if $|\mathfrak{C}_k| \neq 0$, and $\Psi_k(t, \xi, z) = 0$, otherwise, where $s_{(k|\ell)}$ is the Schur function associated with the partition $(k|\ell)$ in Frobenius' notation, and

$$\Theta_{k,q}(t, \xi, x) = \sum_{r=0}^q \frac{1}{(q-r)!} \left(-\frac{1}{\sqrt{2t}} \right)^{q-r} H_{q-r} \left(\frac{c_k - x}{\sqrt{2t}} \right) h_r \left(\left(\frac{1}{u - c_k} \right)_{u \in \xi - \mathfrak{C}_k} \right) \quad (4.10)$$

with the Hermite polynomials $H_k, k \in \mathbb{N}_0$, and with the complete symmetric functions $h_k, k \in \mathbb{N}_0$.

Lemma 4.5 *Suppose that $\xi \in \mathfrak{Y}_0$. Then for $k \in \mathbb{Z}, t \geq 0, x \in \mathbb{R}, z \in \mathbb{C}$*

$$\int_{\mathbb{R}} \mathfrak{C}_k(dx') e^{-(x'-x)^2/(2t)} \Phi(\xi, x', z) = e^{-(c_k-x)^2/(2t)} \Psi_k(t, \xi, z, x).$$

Proof. From definitions of $\mathfrak{C}_k, k \in \mathbb{Z}$ and Φ , we have

$$\begin{aligned} &\int_{\mathbb{R}} \mathfrak{C}_k(dx') e^{-(x'-x)^2/(2t)} \Phi(\xi, x', z) \\ &= \int_{\mathbb{R}} \mathfrak{C}_k(dx') e^{-(x'-x)^2/(2t)} \prod_{u \in \xi - \mathfrak{C}_k} \frac{z - u}{x' - u} \prod_{v \in \mathfrak{C}_k - \delta_{x'}} \frac{z - v}{x' - v} \\ &= e^{-(c_k-x)^2/(2t)} \int_{\mathbb{R}} \mathfrak{C}_k(dx') e^{-(x'-c_k)(x'+c_k-2x)/(2t)} \\ &\quad \times \prod_{u \in \xi - \mathfrak{C}_k} \frac{(z - c_k) - (u - c_k)}{(x' - c_k) - (u - c_k)} \prod_{v \in \mathfrak{C}_k - \delta_{x'}} \frac{(z - c_k) - (v - c_k)}{(x' - c_k) - (v - c_k)} \\ &= e^{-(c_k-x)^2/(2t)} \sum_{j=1}^{|\mathfrak{C}_k|} \psi_k(t, \xi, v_{kj} - c_k, z, x) \frac{a_{\delta}(\mathbf{v}_k - c_k; j; z - c_k)}{a_{\delta}(\mathbf{v}_k - c_k)}, \end{aligned}$$

where

$$a_{\delta}(\mathbf{x}_m; j; y) = a_{\delta}(x_1, \dots, x_{j-1}, y, x_{j+1}, \dots, x_m),$$

and

$$\psi_k(t, \xi, x', z, x) = \Phi(\xi - \mathfrak{C}_k, x' + c_k, z) \exp \left(-\frac{2(c_k - x)x' + x'^2}{2t} \right).$$

Now we introduce $\tilde{\Theta}_{k,q}$'s as the coefficients of the expansion

$$\psi_k(t, \xi, x', z, x) = \sum_{q \in \mathbb{N}_0} \tilde{\Theta}_{k,q}(t, \xi, z, x) x'^q.$$

Then we have

$$\begin{aligned} & \frac{1}{a_\delta(\mathbf{v}_k - c_k)} \sum_{j=1}^{|\mathfrak{C}_k|} \psi_k(t, \xi, v_{kj} - c_k, z, x) a_\delta(\mathbf{v}_k - c_k; j; z - c_k) \\ &= \frac{1}{a_\delta(\mathbf{v}_k - c_k)} \sum_{\ell=1}^{|\mathfrak{C}_k|} (z - c_k)^{\ell-1} (-1)^{|\mathfrak{C}_k|-\ell-1} \\ & \quad \times \det \begin{pmatrix} \psi_k(t, \xi, v_{k1} - c_k, z, x) & \psi_k(t, \xi, v_{k2} - c_k, z, x) & \cdots & \psi_k(t, \xi, v_{k|\mathfrak{C}_k|} - c_k, z, x) \\ (v_{k1} - c_k)^{|\mathfrak{C}_k|-1} & (v_{k2} - c_k)^{|\mathfrak{C}_k|-1} & \cdots & (v_{k|\mathfrak{C}_k|} - c_k)^{|\mathfrak{C}_k|-1} \\ \cdots & \cdots & \cdots & \cdots \\ (v_{k1} - c_k)^{\ell+1} & (v_{k2} - c_k)^{\ell+1} & \cdots & (v_{k|\mathfrak{C}_k|} - c_k)^{\ell+1} \\ (v_{k1} - c_k)^{\ell-1} & (v_{k2} - c_k)^{\ell-1} & \cdots & (v_{k|\mathfrak{C}_k|} - c_k)^{\ell-1} \\ \cdots & \cdots & \cdots & \cdots \\ v_{k1} - c_k & v_{k2} - c_k & \cdots & v_{k|\mathfrak{C}_k|} - c_k \\ 1 & 1 & \cdots & 1 \end{pmatrix} \\ &= \sum_{\ell=1}^{|\mathfrak{C}_k|} (z - c_k)^{\ell-1} (-1)^{|\mathfrak{C}_k|-\ell-1} \\ & \quad \times \left\{ \tilde{\Theta}_{k,\ell}(t, \xi, z, x) + \sum_{q=|\mathfrak{C}_k|}^{\infty} \tilde{\Theta}_{k,q}(t, \xi, z, x) s_{(q-|\mathfrak{C}_k|)(|\mathfrak{C}_k|-\ell-1)}(\mathbf{v}_k - c_k) \right\}. \end{aligned}$$

Then, to prove the lemma, it is enough to show the equality

$$\tilde{\Theta}_{k,q}(t, \xi, z, x) = \Phi(\xi - \mathfrak{C}_k, c_k, z) \Theta_{k,q}(t, \xi, x), \quad t \geq 0, \xi \in \mathfrak{Y}_0, z \in \mathbb{C}, x \in \mathbb{R}, \quad (4.11)$$

for $|\mathfrak{C}_k| \neq 0$. From the formula (3.3), we have

$$\begin{aligned} \Phi(\xi - \mathfrak{C}_k, x' + c_k, z) &= \prod_{u \in \xi - \mathfrak{C}_k} \frac{z - u}{x' - (u - c_k)} \\ &= \prod_{u \in \xi - \mathfrak{C}_k} \frac{u - z}{u - c_k} \prod_{u \in \xi - \mathfrak{C}_k} \frac{1}{1 - x'/(u - c_k)} \\ &= \prod_{u \in \xi - \mathfrak{C}_k} \frac{u - z}{u - c_k} \sum_{r \in \mathbb{N}_0} h_r \left(\left(\frac{1}{u - c_k} \right)_{u \in \xi - \mathfrak{C}_k} \right) x'^r \\ &= \Phi(\xi - \mathfrak{C}_k, c_k, z) \sum_{r \in \mathbb{N}_0} h_r \left(\left(\frac{1}{u - c_k} \right)_{u \in \xi - \mathfrak{C}_k} \right) x'^r. \end{aligned} \quad (4.12)$$

By the formula (3.16), we have

$$\exp\left(-\frac{2(c_k - x)x' + x'^2}{2t}\right) = \sum_{k \in \mathbb{N}_0} \frac{1}{k!} \left(-\frac{x'}{\sqrt{2t}}\right)^k H_k\left(\frac{c_k - x}{\sqrt{2t}}\right). \quad (4.13)$$

Combining (4.12) and (4.13), we have

$$\begin{aligned} & \psi_k(t, \xi, x', z, x) \\ &= \Phi(\xi - \mathfrak{C}_k, c_k, z) \sum_{r \in \mathbb{N}_0} h_r \left(\left(\frac{1}{u - c_k} \right)_{u \in \xi - \mathfrak{C}_k} \right) x'^r \\ & \quad \times \sum_{k \in \mathbb{N}_0} \frac{1}{k!} \left(-\frac{x'}{\sqrt{2t}} \right)^k H_k \left(\frac{c_k - x}{\sqrt{2t}} \right) \\ &= \Phi(\xi - \mathfrak{C}_k, c_k, z) \sum_{q \in \mathbb{N}_0} x'^q \sum_{r=0}^q \frac{1}{(q-r)!} \left(-\frac{1}{\sqrt{2t}} \right)^{q-r} \\ & \quad \times H_{q-r} \left(\frac{c_k - x}{\sqrt{2t}} \right) h_r \left(\left(\frac{1}{u - c_k} \right)_{u \in \xi - \mathfrak{C}_k} \right). \end{aligned}$$

Then, by definition (4.10), (4.11) is proved. ■

Lemma 4.6 Assume that (C.3) holds with some $\kappa \in (1/2, 1)$ and $m \in \mathbb{N}$.

(i) Suppose that $\alpha \in (1/\kappa, 2)$. Then there exists $C_4(\kappa, m, \alpha) > 0$ such that

$$M_\alpha(\tau_{-a}(\xi - \mathfrak{C}^a)) \leq C_4(\kappa, m)(|a| \vee 1)^{(1-\kappa)/\kappa} \quad \forall a \in \text{supp } \xi, \quad (4.14)$$

and (C.2) (i) holds, that is, there exists $C_1 = C_1(\alpha, \xi)$ such that

$$M_\alpha(\xi) \leq C_1. \quad (4.15)$$

(ii) Suppose that $\beta \in (0, 2\kappa - 1)$. Then $\xi - \mathfrak{C}^a - \widehat{\mathfrak{C}^a}$ satisfies (C.2) (ii) $\forall a \in \text{supp } \xi$, where $\widehat{\mathfrak{C}^a} = \mathfrak{C}_{-k}$ in case $\mathfrak{C}^a = \mathfrak{C}_k$. That is, there exists $C_2(\kappa, m) > 0$ such that

$$M_1\left(\tau_{-a^2}(\xi - \mathfrak{C}^a - \widehat{\mathfrak{C}^a})^{(2)}\right) \leq C_2(\kappa, m)(|a| \vee 1)^{-\beta} \quad \forall a \in \text{supp } \xi. \quad (4.16)$$

Proof. By simple calculations we see that there exists a positive constant $C(\kappa)$ such that

$$M_\alpha(\tau_{-a}\eta^\kappa) \leq C(\kappa)(|a| \vee 1)^{(1-\kappa)/\kappa} \quad \forall a \in \text{supp } \eta^\kappa. \quad (4.17)$$

Suppose that $\mathfrak{C}^a = \mathfrak{C}_k$, $k \in \mathbb{Z}$. Then $\xi - \mathfrak{C}^a = \xi \cap [\bar{b}_{k-1}, \underline{b}_k]^c$. We divide the set $[\bar{b}_{k-1}, \underline{b}_k]^c$ into the following four sets:

$$\begin{aligned} A_1 &= \left(-\infty, g^\kappa(k-2)\right], & A_2 &= \left(g^\kappa(k-2), \bar{b}_{k-1}\right), \\ A_3 &= \left(\underline{b}_k, g^\kappa(k+2)\right), & A_4 &= \left[g^\kappa(k+2), -\infty\right). \end{aligned}$$

Then we have

$$\left(\int_{\mathbb{R}} \frac{(\xi - \mathfrak{C}^a)(dx)}{|x - a|^\alpha} \right)^{1/\alpha} \leq \sum_{j=1}^4 \left(\int_{A_j} \frac{\xi(dx)}{|x - a|^\alpha} \right)^{1/\alpha}.$$

From (4.7) and (4.8), we have

$$\begin{aligned} \int_{A_1} \frac{\xi(dx)}{|x - a|^\alpha} &\leq m \sum_{-\infty < \ell \leq k-2} \frac{1}{|g^\kappa(\ell) - g^\kappa(k-1)|^\alpha}, \\ \int_{A_2} \frac{\xi(dx)}{|x - a|^\alpha} &\leq 2m \left(\frac{1}{\varepsilon_{k-1}} \right)^\alpha, \\ \int_{A_3} \frac{\xi(dx)}{|x - a|^\alpha} &\leq 2m \left(\frac{1}{\varepsilon_k} \right)^\alpha, \\ \int_{A_4} \frac{\xi(dx)}{|x - a|^\alpha} &\leq m \sum_{k+2 \leq \ell < \infty} \frac{1}{|g^\kappa(\ell) - g^\kappa(k+1)|^\alpha}. \end{aligned}$$

Combining these estimates with (4.17), we have

$$\left(\int_{\mathbb{R}} \frac{(\xi - \mathfrak{C}^a)(dx)}{|x - a|^\alpha} \right)^{1/\alpha} \leq \mathcal{O}\left((|g^\kappa(k-1)| \vee |g^\kappa(k+1)| \vee 1)^{(1-\kappa)/\kappa}\right), \quad |k| \rightarrow \infty.$$

Since $\max_{k-1 \leq j \leq k+1} |g^\kappa(j)| \leq 2(|a| \vee 1)$, we obtain (4.14). The estimate (4.15) is derived from (4.14) with $a = 0$ and $\mathfrak{C}^a = \mathfrak{C}_0$, and the fact that $M_\alpha(\mathfrak{C}_0) < \infty$. Noting that $(\xi - \mathfrak{C}^a - \widehat{\mathfrak{C}^a})^{(2)}$ satisfies **(C.3)** with 2κ and $2m$, we obtain (4.16) by a similar argument given above to show (4.14). This completes the proof. ■

Lemma 4.7 *Let $\alpha \in (1, 2)$ and $|a| \geq 1$. Assume that **(C.1)** and the condition*

$$M_\alpha(\tau_{-a}\xi) \leq C_5|a|^\gamma \tag{4.18}$$

with some $\gamma > 0$ and $C_5 > 0$ are satisfied. Then there exists $C_6 = C_6(\alpha, \beta, C_1, C_5) > 0$ such that

$$|M(\tau_{-a}\xi) - M(\xi)| \leq C_6|a|^{\delta_1},$$

where $\delta_1 = \alpha(1 + \gamma) - 1$.

Proof. From Lemma 4.3 and the fact that $M_1(\tau_{-a}\xi, L)$ is increasing in L , we see that

$$\max_{0 \leq L \leq L_0} M_1(\tau_{-a}\xi, L) = M_1(\tau_{-a}\xi, L_0) \leq (2M_\alpha(\tau_{-a}\xi))^{\alpha\delta_1/(\delta_1-\alpha+1)} \leq C|a|^{\delta_1}$$

from (4.18) with a constant $C > 0$. Combining this estimate with Lemma 4.3, we have

$$M_1(\tau_{-a}\xi, L) \leq C|a|^{\delta_1} \vee L^{\delta_1}. \tag{4.19}$$

We assume $a \neq 0$. By the definitions of $M(\xi)$ and $M(\tau_{-a}\xi)$,

$$|M(\tau_{-a}\xi) - M(\xi)| \leq \frac{1 + \xi(\{0\})}{|a|} + |a| \int_{\{a,0\}^c} \frac{\xi(dx)}{|x(x-a)|}.$$

We divide the set $\{a,0\}^c$ into the three disjoint subsets $\{x : 0 < |x| < 2|a|, 2|a-x| > |a|\}$, $\{x : |x| \geq 2|a|\}$ and $\{x : 0 < |x| < 2|a|, 0 < 2|a-x| \leq |a|\}$. By simple calculation, we see

$$\int_{0 < |x| < 2|a|, 2|a-x| > |a|} \frac{\xi(dx)}{|x(x-a)|} \leq \frac{2}{|a|} \int_{0 < |x| < 2|a|} \frac{\xi(dx)}{|x|} = \frac{2}{|a|} M_1(\xi, 2|a|).$$

Since $|x-a| \geq |x| - |a| \geq |x|/2$, if $|x| \geq 2|a|$,

$$\int_{|x| \geq 2|a|} \frac{\xi(dx)}{|x(x-a)|} \leq 2 \int_{|x| \geq 2|a|} \frac{\xi(dx)}{|x|^2} \leq 2^{\alpha-1} M_\alpha(\xi)^\alpha |a|^{\alpha-2}.$$

Since $|x| \geq |a| - |a-x| \geq |a|/2$, if $2|a-x| \leq |a|$,

$$\int_{0 < |x| < 2|a|, 0 < 2|a-x| \leq |a|} \frac{\xi(dx)}{|x(x-a)|} \leq \frac{2}{|a|} \int_{0 < 2|a-x| \leq |a|} \frac{\xi(dx)}{|x-a|} = \frac{2}{|a|} M_1\left(\tau_{-a}\xi, \frac{|a|}{2}\right).$$

Combining the above estimates with the fact $|a|^{-1} \leq 1$, we have

$$|M(\tau_{-a}\xi) - M(\xi)| \leq 2^{\alpha-1} M_\alpha(\xi)^\alpha |a|^{\alpha-1} + 2M_1(\xi, 2|a|) + 2M_1\left(\tau_{-a}\xi, \frac{|a|}{2}\right) + 2.$$

Then the lemma is derived from (4.18) and (4.19). ■

The following is a key lemma to prove Theorem 2.4.

Lemma 4.8 *Let $t \geq 0, x \in \mathbb{R}$, $\xi \in \mathfrak{Y}_m^\kappa \subset \mathfrak{Y}$ with $\kappa \in (1/2, 1)$ and $m \in \mathbb{N}$. Then for any $\theta \in (3 - 2\kappa, 2)$ there exist positive constants $C_7 = C_7(t, \kappa, C_0, x)$ and $\widehat{C}_7 = \widehat{C}_7(t, \kappa, m, \theta, C_0, x)$ such that*

$$|\Psi_k(t, \xi, iy, x)| \leq \widehat{C}_7 \exp \left[C_7 \left\{ |y|^\theta + |c_k|^\theta \right\} \right], \quad \forall y \in \mathbb{R}, \forall k \in \mathbb{Z}.$$

Proof. We note the equality

$$\Phi(\xi - \mathfrak{C}_k, c_k, iy) = \Phi(\xi - \mathfrak{C}_k - \mathfrak{C}_{-k}, c_k, iy) \Phi(\mathfrak{C}_{-k}, c_k, iy).$$

Let $\beta \in (0, 2\kappa - 1)$ and $\alpha = (1/\kappa, 2)$. By virtue of Lemma 4.6, we can apply Lemma 4.4 for $\xi - \mathfrak{C}_k - \mathfrak{C}_{-k}$ and see that there exist positive constant C_3 and $\theta \in (3 - 2\kappa, 2)$ such that

$$|\Phi(\xi - \mathfrak{C}_k - \mathfrak{C}_{-k}, c_k, iy)| \leq \exp \left[C_3 \left\{ |y|^\theta + |c_k|^\theta \right\} \right], \quad y \in \mathbb{R}, k \in \mathbb{Z}.$$

Here we used the fact that $3 - 2\kappa > 1/\kappa$ for $\kappa \in (1/2, 1)$. Since $\Phi(\mathfrak{C}_{-k}, c_k, iy)$ is a polynomial function of y , we have

$$|\Phi(\xi - \mathfrak{C}_k, c_k, iy)| \leq \widehat{C}_3 \exp \left[C_3 \left\{ |y|^\theta + |c_k|^\theta \right\} \right], \quad y \in \mathbb{R}, \quad k \in \mathbb{Z},$$

for some $\widehat{C}_3 > 0$. Hence, from the definition (4.9) of $\Psi_k(t, \xi, z, x)$, to prove the lemma it is enough to show the following estimates: for any $\ell = 1, 2, \dots, |\mathfrak{C}_k|$,

$$|(z - c_k)^{\ell-1}| = \mathcal{O}(|z|^{|\mathfrak{C}_k|} \vee |c_k|^{|\mathfrak{C}_k|}), \quad |k| \rightarrow \infty, \quad |z| \rightarrow \infty, \quad (4.20)$$

$$|\Theta_{k,\ell}(t, \xi, x)| = \mathcal{O}(|c_k|^\ell), \quad |k| \rightarrow \infty, \quad (4.21)$$

$$\sum_{q=|\mathfrak{C}_k|}^{\infty} \Theta_{k,q}(t, \xi, x) s_{(q-|\mathfrak{C}_k|)|\mathfrak{C}_k|-\ell-1}(\mathbf{v}_k - c_k) \leq \exp \left[C(|c_k|^{\theta'} \vee 1) \right], \quad k \in \mathbb{Z}, \quad (4.22)$$

with some $C = C(t, x) > 0$ and $\theta' < \theta$. Since (4.20) and (4.21) can be confirmed easily, here we show only the proof of (4.22). Since $|v_{k,\ell} - c_k| \leq \Delta_k$, $1 \leq \ell \leq |\mathfrak{C}_k|$, from the fact (3.1)

$$s_{(q-|\mathfrak{C}_k|)|\mathfrak{C}_k|-\ell-1}(\mathbf{v}_k - c_k) \leq \binom{q-\ell-1}{|\mathfrak{C}_k|-\ell-1} \binom{q}{\ell} \Delta_k^q \leq q^{|\mathfrak{C}_k|} \Delta_k^q, \quad q \in \mathbb{N}.$$

Put $\overline{\Delta}_k = \Delta_k + (\varepsilon_{k-1} \wedge \varepsilon_k)/2$, and remind that $\overline{\Delta}_k = \mathcal{O}(c_k^{(\kappa-1)/\kappa})$, $|k| \rightarrow \infty$. Then we have

$$s_{(q-|\mathfrak{C}_k|)|\mathfrak{C}_k|-\ell-1}(\mathbf{v}_k - c_k) \leq C' \overline{\Delta}_k^q, \quad k \in \mathbb{Z}, \quad q \in \mathbb{N},$$

with some positive constant $C' > 0$. Then

$$\begin{aligned} & \Theta_{k,q}(t, \xi, x) s_{(q-|\mathfrak{C}_k|)|\mathfrak{C}_k|-\ell-1}(\mathbf{v}_k - c_k) \\ & \leq C' \sum_{r=0}^q \frac{1}{(q-r)!} \left(\frac{\overline{\Delta}_k}{\sqrt{2t}} \right)^{q-r} \left| H_{q-r} \left(\frac{c_k - x}{\sqrt{2t}} \right) \right| \overline{\Delta}_k^r \left| h_r \left(\left(\frac{1}{u - c_k} \right)_{u \in \xi - \mathfrak{C}_k} \right) \right|, \end{aligned}$$

and thus

$$\begin{aligned} & \sum_{q=|\mathfrak{C}_k|}^{\infty} \Theta_{k,q}(t, \xi, x) s_{(q-|\mathfrak{C}_k|)|\mathfrak{C}_k|-\ell-1}(\mathbf{v}_k - c_k) \\ & \leq C' \sum_{q \in \mathbb{N}_0} \frac{1}{q!} \left(\frac{\overline{\Delta}_k}{\sqrt{2t}} \right)^q \left| H_q \left(\frac{c_k - x}{\sqrt{2t}} \right) \right| \left| \sum_{r \in \mathbb{N}_0} \overline{\Delta}_k^r \left| h_r \left(\left(\frac{1}{u - c_k} \right)_{u \in \xi - \mathfrak{C}_k} \right) \right| \right|. \end{aligned}$$

Since

$$\left| \frac{d^k}{dz^k} e^{2zy - z^2} \right|_{z=0} \leq \frac{d^k}{dz^k} e^{2z|y| + z^2} \Big|_{z=0}, \quad k \in \mathbb{N}, \quad y \in \mathbb{R},$$

we obtain from (4.13)

$$\begin{aligned} & \sum_{q \in \mathbb{N}_0} \frac{1}{q!} \left(\frac{\bar{\Delta}_k}{\sqrt{2t}} \right)^q \left| H_q \left(\frac{c_k - x}{\sqrt{2t}} \right) \right| \\ & \leq \exp \left(\frac{2\bar{\Delta}_k(c_k - x) + \bar{\Delta}_k^2}{2t} \right) = \mathcal{O} \left(\exp \left(\tilde{C} c_k^{1+(\kappa-1)/\kappa} \right) \right), \end{aligned} \quad (4.23)$$

$|k| \rightarrow \infty$, with a constant $\tilde{C} = \tilde{C}(t, x)$. And if $(\xi - \mathfrak{C}_k)(u) \geq 1$, then $|u - c_k| \geq \Delta_k + \varepsilon_{k-1} \wedge \varepsilon_k$ and

$$\frac{1}{1 - \bar{\Delta}_k/|u - c_k|} \leq Cm$$

with a positive constant C . Hence from (3.2)

$$\begin{aligned} & \sum_{r \in \mathbb{N}_0} \bar{\Delta}_k^r \left| h_r \left(\left(\frac{1}{u - c_k} \right)_{u \in \xi - \mathfrak{C}_k} \right) \right| \\ & \leq \exp \left\{ \left| M(\tau_{-c_k}(\xi - \mathfrak{C}_k)) \right| \bar{\Delta}_k + Cm \bar{\Delta}_k^2 M_2(\tau_{-c_k}(\xi - \mathfrak{C}_k))^2 \right\}. \end{aligned} \quad (4.24)$$

Using Lemmas 4.6 and 4.7, we see that

$$\left| M(\tau_{-c_k}(\xi - \mathfrak{C}_k)) \right| \bar{\Delta}_k = \mathcal{O} \left(|c_k|^{\delta_1 + (\kappa-1)/\kappa} \right), \quad |k| \rightarrow \infty,$$

with any $\delta_1 > \{1 + (1 - \kappa)/\kappa\}/\kappa - 1 = 1/\kappa^2 - 1$, and

$$\bar{\Delta}_k^2 M_2(\tau_{-c_k}(\xi - \mathfrak{C}_k))^2 = \mathcal{O} \left(|c_k|^{\alpha(1-\kappa)/\kappa} \bar{\Delta}_k^\alpha \right) = \mathcal{O}(1), \quad |k| \rightarrow \infty.$$

Since $1/\kappa^2 - 1 + (\kappa - 1)/\kappa + 1 + (\kappa - 1)/\kappa = 1/\kappa^2 + 2(\kappa - 1)/\kappa < 3 - 2\kappa$, for $\kappa \in (1/2, 1)$, (4.22) is derived from (4.23) and (4.24). This completes the proof. ■

Proof of Theorem 2.4. (i) By Lemmas 4.5 and 4.8, if $\xi \in \mathfrak{Y}_0$,

$$\begin{aligned} & \int_{\mathbb{R}} \xi(dx') p(s, x|x') \int_{\mathbb{R}} dy' p(t, -iy|y') \Phi(\xi, x', iy) \\ & = \sum_{k \in \mathbb{Z}} p(s, x|c_k) \int_{\mathbb{R}} dy' p(t, -iy|y') \Psi_k(t, \xi, iy', x). \end{aligned} \quad (4.25)$$

Since this equality holds even if we replace ξ by $\xi \cap [-L, L]$ for any $L > 0$,

$$\begin{aligned} & \lim_{L \rightarrow \infty} \mathbb{K}^{\xi \cap [-L, L]}(s, x; t, y) + \mathbf{1}(s > t) p(s - t, x|y) \\ & = \lim_{L \rightarrow \infty} \int_{\mathbb{R}} \xi \cap [-L, L](dx') p(s, x|x') \int_{\mathbb{R}} dy' p(t, -iy|y') \Phi(\xi \cap [-L, L], x', iy) \\ & = \lim_{L \rightarrow \infty} \sum_{k \in \mathbb{Z}} p(s, x|c_k) \int_{\mathbb{R}} dy' p(t, -iy|y') \Psi_k(t, \xi \cap [-L, L], iy', x). \end{aligned}$$

By Lemma 4.8, we can apply Lebesgue's convergence theorem to show that the limit is

$$\sum_{k \in \mathbb{Z}} p(s, x | c_k) \int_{\mathbb{R}} dy' p(t, -iy|y') \Psi_k(t, \xi, iy', x).$$

We can repeat the argument in the proof of Theorem 2.2 given at the end of Section 4.2. Then if $\xi \in \mathfrak{Y}_0$, $(\mathbb{P}_\xi, \Xi(t), t \in [0, \infty))$ is well-defined with the correlation kernel

$$\begin{aligned} \mathbb{K}^\xi(s, x; t, y) &= \sum_{k \in \mathbb{Z}} p(s, x | c_k) \int_{\mathbb{R}} dy' p(t, -iy|y') \Psi_k(t, \xi, iy', x) \\ &\quad - \mathbf{1}(s > t) p(s - t, x | y). \end{aligned} \quad (4.26)$$

It is equal to (2.3) of Theorem 2.2 by the equality (4.25). When $\xi \in \mathfrak{Y} \setminus \mathfrak{Y}_0$, (4.25) is not valid. For any $L > 0$, however, the equality

$$\begin{aligned} &\mathbb{K}^{\xi \cap [-L, L]}(s, x; t, y) + \mathbf{1}(s > t) p(s - t, x | y) \\ &= \sum_{k \in \mathbb{Z}} p(s, x | c_k) \int_{\mathbb{R}} dy' p(t, -iy|y') \Psi_k(t, \xi \cap [-L, L], iy', x) \end{aligned}$$

holds by the continuity with respect to the initial configuration for Dyson's model with finite particles. Then, again by Lemma 4.8 with Lebesgue's convergence theorem, we will obtain the result (4.26).

(ii) By the fact (2.7) and the definition of Ψ_k , we see that for any $k \in \mathbb{N}$, $t \geq 0$, and $x, y' \in \mathbb{R}$

$$\lim_{n \rightarrow \infty} \Psi_k(t, \xi_n, iy', x) = \Psi_k(t, \xi, iy', x).$$

By using Lemma 4.8 we see that, for fixed $t \geq 0, x \in \mathbb{R}$, there exist $\theta \in (1, 2)$, $C_7 = C_7(t, x) > 0$, and $\widehat{C}_7 = \widehat{C}_7(t, x) > 0$ such that

$$|\Psi_k(t, \xi_n, iy', x)| \leq \widehat{C}_7 \exp \left[C_7 \left\{ |y'|^\theta + |c_k|^\theta \right\} \right], \quad k \in \mathbb{Z}, y' \in \mathbb{R}, n \in \mathbb{N}.$$

Therefore, by applying Lebesgue's convergence theorem, we obtain the theorem. ■

Acknowledgments. The present authors would like to thank T. Shirai and H. Spohn for useful comments on the manuscript. A part of the present work was done during the participation of M.K. in the ESI program "Combinatorics and Statistical Physics" (March and May in 2008). M.K. expresses his gratitude for hospitality of the Erwin Schrödinger Institute (ESI) in Vienna and for well-organization of the program by M. Drmota and C. Krattenthaler. M.K. is supported in part by the Grant-in-Aid for Scientific Research (C) (No.21540397) of Japan Society for the Promotion of Science. H.T. is supported in part by the Grant-in-Aid for Scientific Research (KIBAN-C, No.19540114) of Japan Society for the Promotion of Science.

References

- [1] Andrews, G. E., Askey, R., Roy, R.: Special functions. Cambridge: Cambridge University Press, 1999
- [2] Bleher, P. M., Kuijlaars, A. B.: Integral representations for multiple Hermite and multiple Laguerre polynomials. *Ann. Inst. Fourier.* **55**, 2001-2014 (2005)
- [3] Dyson, F. J. : A Brownian-motion model for the eigenvalues of a random matrix. *J. Math. Phys.* **3**, 1191-1198 (1962)
- [4] Eynard, B., Mehta, M. L. : Matrices coupled in a chain: I. Eigenvalue correlations. *J. Phys. A* **31**, 4449-4456 (1998)
- [5] Fritz, J.: Gradient dynamics of infinite point systems. *Ann. Probab.* **15**, 478-514 (1987)
- [6] Grabiner, D. J.: Brownian motion in a Weyl chamber, non-colliding particles, and random matrices. *Ann. Inst. Henri Poincaré, Probab. Stat.* **35**, 177-204 (1999)
- [7] Imamura, T., Sasamoto, T.: Polynuclear growth model with external source and random matrix model with deterministic source. *Phys. Rev. E* **71**, 041606/1-12 (2005)
- [8] Ismail, M. E. H.: Classical and Quantum Orthogonal Polynomials in One Variable. Cambridge: Cambridge University Press, 2005
- [9] Karlin, S., McGregor, J. : Coincidence probabilities. *Pacific J. Math.* **9**, 1141-1164 (1959)
- [10] Katori, M., Tanemura, H.: Noncolliding Brownian motion and determinantal processes. *J. Stat. Phys.* **129**, 1233-1277 (2007)
- [11] Katori, M., Tanemura, H.: in preparation
- [12] Levin, B. Ya.: Lectures on Entire Functions. Translations of Mathematical Monographs, **150**, Providence R. I.: Amer. Math. Soc., 1996
- [13] Macdonald, I. G. : Symmetric Functions and Hall Polynomials. 2nd edition, Oxford: Oxford Univ. Press, 1995
- [14] Mehta, M. L. : Random Matrices. 3rd edition, Amsterdam: Elsevier, 2004
- [15] Nagao, T., Forrester, P. J. : Multilevel dynamical correlation functions for Dyson's Brownian motion model of random matrices. *Phys. Lett.* **A247**, 42-46 (1998)
- [16] Osada, H. : Dirichlet form approach to infinite-dimensional Wiener processes with singular interactions. *Commun. Math. Phys.* **176**, 117-131 (1996)
- [17] Osada, H. : Interacting Brownian motions in infinite dimensions with logarithmic interaction potentials. [arXiv:math.PR/0902.3561](https://arxiv.org/abs/math.PR/0902.3561)

- [18] Shirai, T., Takahashi, Y.: Random point fields associated with certain Fredholm determinants I: fermion, Poisson and boson point process. *J. Funct. Anal.* **205**, 414-463 (2003)
- [19] Soshnikov, A. : Determinantal random point fields. *Russian Math. Surveys* **55**, 923-975 (2000)
- [20] Spohn, H. : Interacting Brownian particles: a study of Dyson's model. In: *Hydrodynamic Behavior and Interacting Particle Systems*, G. Papanicolaou (ed), IMA Volumes in Mathematics and its Applications, **9**, Berlin: Springer-Verlag, 1987, pp. 151-179
- [21] Tracy, C. A., Widom, H.: Differential equations for Dyson processes. *Commun. Math. Phys.* **252**, 7-41 (2004)